

PhD
1939
r

Robln, I.

BOSTON UNIVERSITY

GRADUATE SCHOOL

Dissertation

PERSEVERATION IN DEMENTIA PRAECOX AND MANIC DEPRESSIVE PSYCHOSES

by

Israel Rabin

(B.S. in Ed., Boston University, 1935; A.M., Boston University, 1936)

submitted in partial fulfilment of the

requirements for the degree of

Doctor of Philosophy

1939

THE UNIVERSITY OF

CHICAGO

LIBRARY

IN THE UNIVERSITY OF CHICAGO

LIBRARY

CHICAGO

THE UNIVERSITY OF CHICAGO

LIBRARY

CHICAGO

LIBRARY

CHICAGO

Ph D
1939
or
copy 1

TABLE OF CONTENTS

	Page
INTRODUCTION	1
I. THE NATURE OF PRESERVATION	3
A. Preservation in everyday life	3
B. Historical sources and methods of preservation	3
C. Preservation and Psychology	10
D. Preservation in Psychology	13
II. GENERAL EXPERIMENTATION ON PRESERVATION	16
A. The Subject	16
B. The Apparatus	16
C. Experimental Work in the various phases	24
D. Some Theoretical Considerations	30
III. CORRELATION WITH PSYCHIC ACTIVITY	33
A. Preservation correlated with the various diagnoses (Psychotics)	33
B. Points of Measurement	41
C. Aim of Present Study	43
IV. EXPERIMENTAL METHOD AND PROCEDURES	45
A. Subjects used	47
B. The Preliminary report	49
C. Tests employed in study	50
1. Memory Tests	50

APPROVED

BY

First Reader


Wayland F. Vaulter

Professor of Psychology

Second Reader

Edwin M. Chamberlin

Professor of Psychology



Digitized by the Internet Archive
in 2016 with funding from
Boston Library Consortium Member Libraries

<https://archive.org/details/perseverationind00rabi>

TABLE OF CONTENTS

	Page
INTRODUCTION	1
I. THE NATURE OF PERSEVERATION	3
A. Perseveration in everyday life	3
B. Historical sources concerning perseveration	3
C. Perseveration and Physiology	10
D. Perseveration in Psychiatry	13
II. GENERAL EXPERIMENTATION ON PERSEVERATION	16
A. The Dutch School	16
B. The English School	19
C. Experimental Work in the United States	24
D. Summary of Experimental Methods	28
E. Some Theoretical Considerations	30
III. EXPERIMENTATION WITH PSYCHOTIC MATERIAL	32
A. Perseveration correlated with the various diagnoses (Psychiatric)	32
B. Points of Disagreement	42
C. Aim of Present Study	45
IV. EXPERIMENTAL METHODS AND PROCEDURE	47
A. Subjects used	47
B. The Preliminary rapport	48
C. Tests employed in Study	50
1. Motor Tests	50

TABLE OF CONTENTS

Page

1	INTRODUCTION
2	I. THE NATURE OF PRESERVATION
3	A. Preservation in everyday life
5	B. Historical sources concerning preservation
10	C. Preservation and Physiology
12	D. Preservation in Psychiatry
16	II. GENERAL EXPERIMENTATION ON PRESERVATION
16	A. The Dutch School
19	B. The English School
24	C. Experimental Work in the United States
26	D. Summary of Experimental Methods
30	E. Some Theoretical Considerations
32	III. EXPERIMENTATION WITH PSYCHIC MATERIAL
	A. Preservation correlated with the various
32	diagnoses (Psychiatrists)
42	B. Points of Disagreement
45	C. Aim of Present Study
47	IV. EXPERIMENTAL METHOD AND PROCEDURE
47	A. Subjects used
48	B. The Preliminary report
50	C. Tests employed in Study
50	1. Motor Tests

	Page
a. Reverse Stroke (Z)	51
b. Stroke test	54
2. Sensory Tests	54
a. Light adaptation	56
b. Electrical sensitivity threshold	57
3. A Test of Rhythmic perseveration	57
D. Critique on Subjects' Reports (Reliability)	60
E. Methods of Scoring	61
V. EXPERIMENTAL RESULTS	65
A. Preliminary Results	65
B. Further Results	66
C. The Test on Rhythmic Perseveration	77
VI. DISCUSSION OF RESULTS	79
A. Diagnoses and P-Scores	79
B. Intercorrelation between the Tests	81
C. Relation between Age and Perseveration	82
D. "Improvement" and P-scores	83
E. Theoretical Considerations	84
VII. CONCLUSIONS	86
A. On the Future	87
BIBLIOGRAPHY	89

51	a. Reverse Stroke (C)
52	1. Stroke Test
53	2. Sensory Tests
55	a. Light adaptation
57	1. Electrical sensitivity threshold
57	3. A Test of Rhythmic Perception
59	1. Critique on Subjects' Reports (Reliability)
61	2. Methods of Scoring
62	V. EXPERIMENTAL RESULTS
63	A. Preliminary Results
65	B. Further Results
67	C. The Test on Rhythmic Perception
69	VI. DISCUSSION OF RESULTS
71	A. Response and P-Response
81	B. Interrelation between the Tests
83	C. Relation between Age and Perception
85	D. "Impulsivity" and P-response
84	E. Theoretical Considerations
86	VII. CONCLUSIONS
87	1. In the Future
89	BIBLIOGRAPHY

LIST OF TABLES

	Page
I. Sensory Perseveration in Normals and Psychotics (Wiersma)	33
II. Jones' Results	33
III. P-Scores for Psychotics and Normals (Stephenson) . .	37
IV. Cattell's Results	40
V. Shipley's Results	42
VI. Mean P-Scores in Cases of Dementia Praecox and Manic Depressive	65
VII. Preliminary Correlations of Tests with Age	66
VIII. Means of P-Scores for Dementia Praecox and Manic Depressive Cases	69
IX. P-Scores of Catatonics and other Praecox cases . .	70
X. P-Scores Comparing Catatonic and Paranoiacs . . .	71
XI. P-Scores of Paranoiacs and "Others"	72
XII. A Comparison between Mania and Depression in regard to Perseveration	74
XIII. Correlations Between the Tests	75
XIV. Test Scores for Younger Age Groups	76
XV. Test Scores for Older Age Groups	76
XVI. P-Scores of Improved Subjects	77
XVII. P-Scores for High and Low Perseverators as Determined by the Test on Rhythmic Perseveration	78

LIST OF TABLES

Page	
33	I. Accuracy of Interpretation in Normal and Depressed (1st Form)
35	II. Jones' Results
37	III. P-Scores for Psychotics and Normals (Stephenson)
40	IV. Cattell's Results
42	V. Shipley's Results
	VI. Mean P-Scores in Cases of Depressive Process and Mania
43	Depressive
45	VII. Preliminary Correlations of Tests with Age
	VIII. Means of P-Scores for Depressive Process and Mania
49	Depressive Cases
50	IX. P-Scores of Depressive and other Process Cases
51	X. P-Scores Compared with Psychotics and Paranoics
52	XI. P-Scores of Paranoics and "Others"
	XII. A Comparison between Mania and Depression in regard to
54	Interpretation
55	XIII. Correlations between the Tests
56	XIV. Test Scores for Younger Groups
58	XV. Test Scores for Older Groups
59	XVI. P-Scores of Depressive Subjects
	XVII. P-Scores for High and Low Interpretation as Determined
61	by the Test of Subject's Interpretation

Introduction

Since the end of the last century, when English psychologists began to turn their attention to the study of individual differences, the field of psychology has been enriched by the study of individual differences. The study of individual differences has been enriched by the study of individual differences. The study of individual differences has been enriched by the study of individual differences.

List of Figures

	Page
Figure 1. Record of the Z-Test	53
Figure 2. Record of the Stroke Test	55
Figure 3. Kymographic Record of the Test of Rhythmic Perseveration ("No Change" Patient)	59
Figure 4. Kymographic Record of the Test of Rhythmic Perseveration (A "Change" Patient)	59

List of Figures

Page

Figure 1. Record of the E-Test	51
Figure 2. Record of the E-Test	52
Figure 3. Rhythmic Record of the Test of Rhythmic Interconversion ("No Change" Patient)	53
Figure 4. Rhythmic Record of the Test of Rhythmic Interconversion (A "Change" Patient)	54

Introduction

Since the end of the last century, when English psychologists (Galton and others) brought to the attention of the psychological world the problems of individual differences, much research has been done. Hundreds and, perhaps, thousands of articles, volumes, and monographs have been published on this subject. The quest for the discovery of laws which may, paradoxically enough, explain these individual differences was predominant.

Soon after the British beginning, American psychology took up the challenge and, headed by Cattell, attempted to get at the basic constituents of mind and the laws governing their variability and difference in degree among different individuals.

The new interest in the investigation of traits introduced a new elementism. For a while the traits as such, abstract as they were, were measured with little regard to their interrelationship with other traits and the total, concrete personality. More recently, however, psychology began to admit the error. It started to consider the measured trait within its context, in its psychological background, which is the unified personality. Thus, our analysis into traits is merely temporary; it presumes and anticipates a post-analytic synthesis.

Concurrent with the movement of individual differences, there arose, also, the new tendency to learn something about the normal personality from the study of psychopathological material (Freud, Gross, Jung, Adler and others). The extremes of behavior, as manifested in psychotics, hold, to some extent, the key to our understanding the behavior of the "normal" individual. The gap between the abnormal and the normal is bridged, it must be confessed, by a "process of logic." But, psychology, as it is at present, will have to resort to these devices, until our experimental

Since the end of the last century, when English psychology (Galton and others) brought to the attention of the psychological world the problem of individual differences, much research has been done. Hundreds and, perhaps, thousands of articles, volumes, and monographs have been published on this subject. The quest for the discovery of laws which may, paradoxically enough, explain these individual differences was predominant. Soon after the British beginning, American psychology took up the challenge and, headed by Cattell, attempted to get at the basic consistencies of mind and the laws governing their variability and difference in degree among different individuals.

The new interest in the investigation of traits introduced a new elementism. For a while the traits as such, abstract as they were, were measured with little regard to their interrelationship with other traits and the total, concrete personality. More recently, however, psychology began to admit the error. It started to consider the measured traits within its context, in the psychological background which is the unified personality. Thus, our analysis into traits is merely temporary; it presumes and anticipates a post-analytic synthesis.

Concurrent with the movement of individual differences, there arose, also, the new tendency to learn something about the normal personality from the study of psychopathological material (Freud, Gross, Jung, Adler and others). The extremes of behavior, as manifested in psychotics, hold, to some extent, the key to our understanding the behavior of the "normal" individual. The gap between the abnormal and the normal is bridged, it must be confessed, by a "process of logic." But, psychology, as it is at present, will have to resort to these devices, until our experimental

techniques and the logic behind them are invulnerable and doubtlessly objective and "scientific." Thus far, as to some extent all sciences do, we are dealing with approximations of the truth.

The two tendencies mentioned above seem to be combined in the present study, which is a study of individual differences regarding perseveration in psychotic personalities. Much of the stimulus in this direction was given by Cameron who says that "Perseveration is a personality characteristic from the further study of which much is to be expected."¹ Moreover, he points towards psychiatry as being in need of research directed in that path, and says that "....there is a considerable need for further work in classifying the issues in connection with their function....it may lead to the elucidation of matters of considerable psychiatric importance."²

¹ D.E.Cameron, Objective, Experimental Psychiatry, 70, 1935.

² Ibid, 73.

techniques and the logic behind them are invaluable and doubtless objective and "scientific." Thus far, as to some extent all sciences do, we are dealing with approximations of the truth.

The two tendencies mentioned above seem to be combined in the present study, which is a study of individual differences regarding perseveration in psychotic personalities. Much of the stimulus in this direction was given by Cameron who says that "Perseveration is a personality characteristic from the further study of which much is to be expected."¹ Moreover, he points towards psychology as being in need of research directed in that path, and says that "...there is a considerable need for further work in classifying the issues in connection with their function....it may lead to the elucidation of matters of considerable psychiatric importance."²

4. PERSEVERATION IN EVERYDAY LIFE

Close and critical observation of everyday life will yield many bits of human behavior that can be classified under the general term of perseveration. To some extent, greater or lesser, it is a general property of all human behavior. And, as many other characteristics, it differs in extent and intensity with different individuals.

The difficulty in getting started on some task, the tendency to continue a certain activity once it is begun, the fact that "ruts in the road," the difficulty caused by habit in leaving one activity or behavioral pattern to another, are all illustrative of what is meant by perseveration.

CHAPTER I.

To bring an analogy from physics, it is a form of mental inertia which, presumably, is at much a property of mental energy, as physical inertia is of physical energy.

THE NATURE OF PERSEVERATION

This inertia, "perseveration" or perseverative tendency, moreover, persists despite conscious efforts on the part of the individual to eliminate it and to rid himself of it.

5. Historical Sources Regarding Perseveration

Like psychology itself, perseveration has a long past but a very short history.¹ Many investigators have presented summaries of the non-experimental literature concerning perseveration and have not only reviewed the psychological writings on the subject, but have also traced back the recognition of the phenomenon and its manifestations through the philosophical literature as far back as Plato.² Since this paper is not mainly

¹ A. C. Goring, History of Experimental Psychology, III, 1928.

² G. S. Cooperman, The Evolution of Man, 303, 1927.

CHAPTER I.

THE NATURE OF PRESERVATION

A. Perseveration in Everyday Life

Close and critical observation of everyday life will yield many bits of human behavior that can be classified under the general term of perseveration. To some extent, greater or lesser, it is a general property of all human behavior. And, as many other characteristics, it differs in extent and intensity with different individuals.

The difficulty in getting started at some task, the tendency to continue a certain activity once it is begun, the tune that "runs in the head," the difficulty caused by shifting from one activity or behavioral pattern to another, are all illustrative of what is meant by perseveration. To bring an analogy from physics---it is a sort of mental inertia which, presumably, is as much a property of mental energy, as physical inertia is of physical energy.

This inertia, "perseveration" or perseverative tendency, moreover, persists despite conscious efforts on the part of the individual to eliminate it and to rid himself of it.

B. Historical Sources Regarding Perseveration

Like psychology itself perseveration has a long past but a very short history.¹ Many investigators have presented summaries of the non-experimental literature concerning perseveration and have not only reviewed the psychological writings on the subject, but have also traced back the recognition of the phenomenon and its manifestations through the philosophical literature as far back as Plato.² Since this paper is not mainly

¹ E.C.Boring, History of Experimental Psychology, iii, 1935.

² C.Spearman, The Abilities of Man, 305, 1927.

A. Persistence in Everyday Life

Close and critical observation of everyday life will yield many bits of human behavior that can be classified under the general term of persistence. To some extent, greater or lesser, it is a general property of all human behavior. And, as many other characteristics, it differs in extent and intensity with different individuals.

The difficulty in getting started at some task, the tendency to continue a certain activity once it is begun, the time that "runs in the head," the difficulty caused by shifting from one activity or behavioral pattern to another, are all illustrative of what is meant by persistence. To bring an analogy from physics---it is a sort of mental inertia which, presumably, is as much a property of mental energy, as physical inertia is of physical energy.

This inertial, "persistence" or perseverative tendency, moreover, persists despite conscious efforts on the part of the individual to eliminate it and to rid himself of it.

B. Historical Sources Regarding Persistence

Like psychology itself persistence has a long past but a very short history.¹ Many investigators have presented summaries of the non-experimental literature concerning persistence and have not only reviewed the psychological writings on the subject, but have also traced back the recognition of the phenomenon and its manifestations through the philosophical literature as far back as Plato.² Since this paper is not mainly

¹ E. E. Spence, History of Experimental Psychology, III, 1936.
² E. E. Spence, The Abilities of Man, 1927.

concerned with the historical basis of the term and its philosophical genealogy, and since such broad historical excursions have not yielded new methods of attack upon the investigation of it, the present review will be confined to the psychiatric and psychological literature since the genesis and introduction of the term "perseveration" as such.

It was in 1894 that Neisser³ applied the term "perseveration" for the first time, in describing the repetitious and continuous activities of certain mental patients. Muller and Pilzecker in connection with their experimental work upon memory observed the tendency of certain material to recur in consciousness spontaneously, without any conscious effort upon the part of the observers. This phenomenon was attributed to the certain "forces" possessed by ideas which recur as a result of the so-called "perseverative tendency."⁴ Thus Foster quotes the above mentioned investigators as follows:⁵

1. After one has been busy for several hours in attentive observation of a phenomenon, such for instance as the movement of a pointer on a scale, the phenomenon may suddenly present itself again, in image, with the distinctness of sensory experience, especially at times when attention is not demanded in other directions (Fechner's phenomena of sensory memory).
2. Impressive experiences sometimes reappear in the form of hallucinations, at least at times when attention is not demanded strongly in other directions.
3. A long continued motor activity sets up in the lower centers a tendency automatically to continue or repeat the activity. After a strong disposition of this sort has been established, the center affected may 'reel off' the activity for a certain length of time, without the necessity of further incitement (Motorische Einstellung).
4. Cases of perseveration, in the sense of the psychiatrists.

³ C. Neisser, "65 Sitzung Des Vereins Ostdeutscher Irren Aerzte," Allg. Zsch. f. Psychiatrie, 1894, 51, 1016.

⁴ G. E. Muller and H. Pilzker, "Studien der Gedächtniss," Zsch. f. Psychol., Erg. 1, 1900.

⁵ W. S. Foster, "On the Perseverative Tendency," American Journal of Psychology, 1904, 25, 393-428.

concerned with the historical basis of the term and its philosophical
 genealogy, and since such broad historical questions have not vitiated the
 methods of attack upon the investigation of it, the present review will be
 confined to the psychiatric and psychological literature since the genesis
 and introduction of the term "perseveration" as such.

It was in 1894 that Kraepelin¹ applied the term "perseveration" for the
 first time, in describing the repetitions and compulsive activities of
 certain mental patients. Miller and Pillsbury² in connection with their
 experimental work upon memory observed the tendency of certain material to
 recur in conscious/unconscious spontaneously, without any conscious effort upon
 the part of the observer. This phenomenon was attributed to the certain
 "forces" possessed by ideas which recur as a result of the so-called
 "perseverative tendency."³ Thus Foster⁴ traces the above mentioned investi-

gations as follows:

1. "After one has been busy for several hours in attentive observation
 of a phenomenon, such for instance as the movement of a pointer on a
 dial, the phenomenon may suddenly present itself again, in image, with
 the distinctness of sensory experience, especially at times when
 attention is not demanded in other directions (Fechner's phenomenon of
 sensory memory)."
2. "Inceptive experiences sometimes reappear in the form of hallucina-
 tions, at least at times when attention is not demanded strongly in
 other directions."
3. "A long continued motor activity sets up in the lower centers a
 tendency automatically to continue or repeat the activity. After a
 strong disposition of this sort has been established, the center affected
 may 'run off' and actually for a certain length of time, without the
 necessity of further incentive (Motorische Erregung)."
4. Cases of perseveration, in the sense of the psychiatrist.

¹ C. Kraepelin, "Das Mania und Depressionen," Leipzig, 1904, 1014.
² E. C. Miller and E. Pillsbury, "Studies in Psychopathology," Psychol. Rev., 1, 1900.
³ W. S. Foster, "On the Perseverative Tendency," American Journal of
 Psychology, 1904, 25, 222-223.

5. Quite against our will, ideas and thought in which we have been deeply interested for some time---melodies, chess games, and the like, may keep coming back into our minds.

6. We often make mistakes, in reading, hearing, speaking or writing words where words had in mind previous to the mistakes seem, by their continuance (Nachklingen), responsible for the disturbance.

7. In experiments on memory, ideas given as reactions in the first part of the series may recur during the latter part of the series without the volition of the observer, and without any evidence of special reproductive tendencies.

8. In spite of instructions not to think of the series, many observers cannot prevent certain syllables of a series which they have learned from coming to mind in the interval between the learning and the test.

9. In experiments by the methods of right associates, not only may a syllable which has caught the observer's attention be given once or oftener as a response in the rest of the series of which it was a part, but it may also recur as a response on a number of later days (habituelle Aushilfssilben).

10. Incorrect responses, in which the syllable spoken is a syllable from the series, but is given in the wrong place, are much more frequent in tests which follow at longer intervals, even though the number of correct responses in the two cases is approximately the same.

11. Series tested after a short interval give shorter reaction times than series tested after long intervals, even though the number of correct replies is approximately equal in the two cases....

12. Mental activity during a period shortly after learning not only disturbs the associative connections (retroactive inhibition), but also brings it about that the syllables are less likely to rise freely into consciousness.

13. Observers differ greatly in respect to the degree in which the above mentioned symptoms of perseverative tendency appear.

The conception of perseveration was considerably broadened and supplemented by the later contribution of Otto Gross,⁶ who drew largely upon his psychopathological clinical experience. However, as he says in the introduction to his monograph,---psychotic material merely accentuates certain qualities and gives us a chance to study the extremes from the

⁶ O. Gross, Die Zerebrale Sekundaerfunktion, 69, 1902.

6. Quite against our will, ideas and thoughts in which we have been heavily interested for some time--melodies, chess games, and the like, may keep coming back into our minds.

7. We often make mistakes, in reading, hearing, speaking or writing words where words had in mind previous to the mistakes were, by their consonance (Wohlklang), responsible for the disturbance.

8. In experiments on memory, ideas given as questions in the first part of the series may recur during the latter part of the series without the volition of the observer, and without any evidence of special reproductive tendencies.

9. In spite of instructions not to think of the series, many observers cannot prevent certain syllables of a series which they have learned from coming to mind in the interval between the learning and the test.

10. In experiments by the methods of right associations, not only may a syllable which has caught the observer's attention be given once or often as a response in the rest of the series of which it was a part, but it may also occur as a response on a number of later days (spontaneous associations).

11. Incorrect responses, in which the syllable spoken is a syllable from the series, but is given in the wrong place, are much more frequent in tests which follow at longer intervals, even though the number of correct responses in the two cases is approximately the same.

12. Series tested after a short interval give shorter reaction times than series tested after long intervals, even though the number of correct replies is approximately equal in the two cases....

13. Mental activity during a period shortly after learning not only disturbs the associative connections (reproductive inhibition), but also brings it about that the syllables are less likely to be freely introduced into connections.

14. Observers differ greatly in respect to the degree in which the above mentioned symptoms of perseverative tendency appear.

The conception of perseveration was considerably broadened and

supplemented by the later contribution of Otto Gross, who drew largely upon his psychopathological clinical experience. However, as he says in the introduction to his monograph, "Psychotic material merely occasions certain questions and gives us a chance to study the answers from the

middle; the "normal" can be understood. His psychopathological investigations point toward an ultimate understanding of certain "Individualitaets-verschiedenheiten" (individual differences).

Mental life is divided by Gross into the primary functions, which are synonymous with regular perception and ideation, and secondary functions, or after-functions. According to him:⁷

In every nervous element the functional stimulation of which means a representation in consciousness, there lingers after the disappearance of that representation from consciousness, an after-function which no more possesses a correlate in consciousness and which has a further directive effect upon the associational activity.

This "Nachfunction" or "Sekundaerfunction," as he later refers to it, is according to Gross, different in different individuals. The strength and intensity of that secondary function is dependent on the type of consciousness possessed by the person in question. At this point the author resorts to a dichotomy which apparently had considerably influenced Jung in his creation of Introversion and Extroversion. Gross seems to classify human consciousnesses into two main types---the "narrow and deep" type and the "broad superficial" type of consciousness. The latter type finds its most extreme and salient expression in the manic condition of insanity in which:⁸

....der Reiz von aussen ist staerker als die Sekundaerfunction frueherer Vorstellungen, das Individuum wendet sich ungehemmt mit seiner Aufmerksamkeit dem ausseren Reizen zu.*

Thus there is the ability to shift quickly from one stimulus to another, from one activity to another without too much interference on

⁷ ibid, 10. (Translation---my own.)

⁸ ibid, 22.

*When....the outward stimulus is stronger than the secondary function of previous ideas, the individual turns his attention, without inhibition, upon this outside stimulus. (Translation---my own.)

middle; the "normal" can be understood. His psychological investigation point toward an intimate understanding of certain "individual differences" (individual differences).

mental life is divided by Gross into the primary functions, which are synchronous with regular perception and ideation, and secondary functions or after-functions. According to him:

In every nervous element the functional stimulation of which means a representation in consciousness, there lingers after the disappearance of that representation from consciousness, an after-function which no more possesses a correlate in consciousness and which has a further directive effect upon the associational activity.

This "Nachwirkung" or "after-function," as he later refers to it, is according to Gross, different in different individuals. The strength and intensity of that secondary function is dependent on the type of consciousness possessed by the person in question. At this point the author resorts to a dichotomy which apparently had considerably influenced him in his creation of introversion and extroversion. Gross seems to classify human consciousness into two main types--the "narrow and deep" type and the "broad superficial" type of consciousness. The latter type finds its most extreme and salient expression in the manic condition of insanity in which:

....der Mann von aussen ist attackiert als die Nachwirkung
 ihrer Vorstellungen, das Individuum wandert sich ungeschwächt mit
 seiner Aufmerksamkeit den ausseren Reizen zu.*

Thus there is the ability to shift quickly from one stimulus to another, from one activity to another without too much interference on

Y 1918, 10. (Transition--my own.)
 S 1918, 22.

*When....the outside stimulus is stronger than the secondary function of previous ideas, the individual turns his attention, without inhibition, upon this outside stimulus. (Transition--my own.)

account of the lag of the secondary function.

The former type, however, does not shift his attention so quickly. In this "narrow and deep" consciousness "Vorstellungen" persist, make many connections and, skipping from subject to subject, from stimulus to stimulus, are prevented by the strong power of an enhanced secondary function. On the pathological side melancholia is an important harbor for that lag and after-function and, thanks to the latter, it is characterized by that monotony due to the existing "Intrapsychic Stereotypy."⁹

As a consequence of this characterization of psychiatric groups, on the basis of the strength of their secondary function, he also comes to the typology among normals, i.e. to perseverators of narrow consciousness and to non-perseverators of broad consciousness. One thing must be reiterated, though stated in Gross' words on an earlier page---the secondary function that persists modifies the mental processes that follow it. It is essential in reasoning and in connecting "Vorstellungen," it lends a certain continuity to the mental process. The manic flight of ideas shows extremely little secondary function in contradistinction to the melancholic and paranoiac, whose excess of the secondary function keeps him dwelling on the same subject.

Thus, while Muller and Pilzecker spoke of ideas having forces that appear suddenly into consciousness as a result of favorable conditions (no other strong forces present), Gross posited a similar notion of a secondary function, characteristic of nervous plasm persisting for a time as an after-function, unconscious, though influential in subsequent mental activity.

Special importance is attached by Jung¹⁰ to the phenomenon which was

⁹ Ibid, 40.

¹⁰ C. Jung, Collected Papers on Analytical Psychology, 1922.

account of the lag of the secondary function.

The former type, however, does not shift his attention so quickly.

In this "narrow and deep" consciousness "Vorstellungsgesamtheit" persists, makes many connections and, skipping from subject to subject, from stimulus to stimulus, are prevented by the strong power of an enhanced secondary function.

On the pathological side melancholia is an important factor for that lag

and after-function and, thanks to the latter it is characterized by that

monotony due to the existing "Intrapsychische Identifizierung".⁹

As a consequence of this characteristic of psychopathic groups, on

the basis of the strength of their secondary function, it also comes to the

typology among normals, i.e. to persistence of narrow consciousness and

to non-persistence of broad consciousness. One thing must be repeated,

though stated in Gross' words on an earlier page--the secondary function

that persists modifies the mental processes that follow it. It is essential

in reasoning and in connecting "Vorstellungen", it leads a certain continuity

to the mental process. The mental flight of ideas shows extremely little

secondary function in contrast to the melancholic and paranoiac,

whose excess of the secondary function keeps him dwelling on the same sub-

ject.

Thus, while Miller and Fiszbein spoke of ideas having forces that

appear suddenly into consciousness as a result of favorable conditions (no

other strong forces present), Gross posited a similar notion of a secondary

function, characteristic of nervous system persistence for a time as an after-

function, unconscious, though influential in subsequent mental activity.

Special importance is attached by Jung¹⁰ to the phenomenon which was

hinted at by Gross and was termed emotional perseveration. From his studies on the association processes Jung observed that "....they (the perseverated words) always contain something of importance for the individual psychology of the test person."¹¹ One might say that association words which reveal a great deal of secondary function also reveal high emotional tone or "complexes" behind them. Even more significant than the above is the close connection between Jung's typology and that of Gross. The latter showed two processes which are widely separated from each other in regard to the secondary function and which serve as a basis for the typology.¹²

Der Steigerung des Sekundaerfunktion entspricht eine Verengung und event. eine Vertiefung, des Herabminderung der Sekundaerfunktion eine Verflachung und event. eine Verbreitung des Bewusstseins.*

Jung¹³ acknowledges his debt to Gross in his outline of the extrovert type which essentially coincides with Gross' broad and shallow consciousnesses, characteristic of the short secondary function of time. The long secondary function, especially in affective life, characteristic of the "deep and narrow" type of Gross and the "Strong relief given in this picture to the inner processes and the emotional life...."¹⁴ reveal the introvert immediately.

It was not strange that many investigations, as a result of psychiatric observations, and the above-mentioned theoretical considerations, were devoted to the exploration and the establishment of the true relationship

¹¹ Ibid, 106.

¹² O. Gross, Die Zerebrale Sekundaerfunktion, 18, 1902.

¹³ C. Jung, Psychological Types, 341, 1926.

¹⁴ Ibid, 346.

*The rise of the secondary function represents a narrowing and, eventually, a deepening and the decrease of the secondary function, a shallowing and eventually a broadening of consciousness. (Translation---my own.)

indicated as by Gross and was termed emotional perversion. From his studies on the association processes Jung observed that "...they (the perverted words) always contain something of importance for the individual psychology of the test person." ¹¹ One might say that association words which reveal a great deal of secondary function also reveal high emotional tone or "over-pleasure" behind them. Even more significant than the above is the close connection between Jung's typology and that of Gross. The latter showed two processes which are widely separated from each other in regard to the secondary function and which serve as a basis for the typology. ¹²

Der Abgleichung des Hauptgehirnfunktion entspricht eine Verengung und event. eine Verflüchtung, des Nebengehirnfunktion eine Verflüchtung und event. eine Verengung des Bewusstseins. Jung ¹³ acknowledges his debt to Gross in his outline of the extroverted type which essentially coincides with Gross' broad and shallow connectionness, characteristic of the short secondary function of time. The long secondary function, especially in affective life, characteristic of the deep and narrow type of Gross and the "strong" trait given in this picture to the inner processes and the emotional life.... ¹⁴ reveal the intimate relationship.

It was not strange that many investigations, as a result of psychobiological observations, and the above-mentioned theoretical considerations, were devoted to the exploration and the establishment of the true relationship.

- 11 Jung, 1903, 1904, 1905.
- 12 C. Gross, Die Gehirnfunktion, 1903.
- 13 C. Jung, Psychologische Typen, 1928.
- 14 Jung, 1903.

*The idea of the secondary function represents a narrowing and eventually a deepening and the duration of the secondary function, a widening and eventually a prolongation of connectionness. (Translation—)

between the extremes, as well as the finer groupings of extroversion and introversion, and the secondary function or the perseverative tendency.

The extremes on the introversion-extroversion continuum were to be found in the psychopathological groups; the melancholics were considered to be on the introvert end of the scale, while the manics were at the other end of extroversion. The less extreme cases were those of normal individuals whose tendencies put them more or less definitely in one category (introvert or extrovert).

No discussion of the historico-theoretical background of perseveration would be complete without mentioning the impetus given by Spearman to the further experimental research in this field. He admits that on the basis of what is known thus far, perseveration is only a "half truth", especially as far as its establishment as a true faculty is concerned; but investigations of it have furnished "....the most lavish of promises for individual psychology."¹⁵

Spearman formulated a law of "lag or inertia" which states that: "Cognitive processes always both begin and cease more gradually than their (apparent) causes."¹⁶ Spearman, in his discussion of perseveration, however, is not quite satisfied with what is known about perseveration, though the experimental evidence quoted by him seems to point towards its establishment as a group factor possessing functional unity. This latter conclusion, as it will be pointed out later (Chapter II), was a bit of premature optimism, since more critical experimentation seemed to shatter its foundations and put under question its right to existence. Of great interest to Spearman

¹⁵ C. Spearman, The Abilities of Man, 306, 1927.

¹⁶ Ibid, 291.

is "....how this second universal factor stands in relation to 'g'."¹⁷
 Since "g" is the general factor of intelligence and "p" (perseveration) is another factor---its degree of inertia; the relation between the two is a cardinal problem worth investigating.

Concluding this section it would be, perhaps, best to state Edwards,¹⁸ summary of the types of perseveration which will restate briefly the main characteristics of the phenomenon.

When a given note or combination of notes is struck there may be a continuance of sensory impressions long after the stimulus has ceased. Secondly, there may be a later spontaneous revival into consciousness of the note or combination of notes without the original stimulus. Thirdly, there may be the continuance of the past experience which hinders the reaction to a present experience, usually related to the former.

Thus, three phenomena are emphasized above, which seem to summarize perseveration in action---(1) continuance, (2) spontaneous recurrence, which is quite akin to continuance, and (3) interference, which is also a result of continuance of sensations or their inertia - "Nachfunction."

C. Perseveration and Physiology

Some of the earlier objections to the term perseveration or perseverative tendency were raised on physiological grounds as well as on psychological ones. Thus Foster states:¹⁹

Not only may the necessity of supposing a spontaneous recurrence of mental contents be questioned, but a spontaneous activity of the nervous system, which seems thereby to be implied, is also made improbable by physiological facts so far known.

¹⁷ ibid, 306.

¹⁸ K.H.R. Edwards, "The Perseverative Tendency," The Journal of Abnormal and Social Psychology, 1933, 28, 198-203.

¹⁹ W.S. Foster, "On the Perseverative Tendency," American Journal of Psychology, 1914, 25, 393-428.

is "...how this second universal factor stands in relation to 'g'." Since 'g' is the general factor of intelligence and 'p' (perseveration) is another factor---the degree of inertia; the relation between the two is a cardinal problem worth investigating.

Considering this section it would be, perhaps, best to state briefly the main summary of the types of perseveration which will restate briefly the main characteristics of the phenomenon.

When a given note or combination of notes is struck there may be a continuance of sensory impressions long after the stimulus has ceased. Secondly, there may be a later spontaneous revival into consciousness of the note or combination of notes without the original stimulus. Thirdly, there may be the continuance of the past experience which hinders the reaction to a present experience, usually related to the former.

Thus, three phenomena are emphasized above, which seem to summarize perseveration in action---(1) continuance, (2) spontaneous recurrence, which is quite akin to continuance, and (3) interference, which is also a result of continuance of sensations or their inertia - "Nachwirkung."

C. Perseveration and Psychology

Some of the earlier objections to the term perseveration or perseverative tendency were raised on physiological grounds as well as on psychological ones. Thus Foster states:

Not only may the necessity of emphasizing a spontaneous recurrence of mental contents be questioned, but a spontaneous activity of the nervous system, which seems thereby to be implied, is also made improbable by physiological facts so far known.

1. Journal of Psychology, 1914, 25, 325-332.
 2. Journal of Psychology, 1914, 25, 325-332.
 3. Journal of Psychology, 1914, 25, 325-332.
 4. Journal of Psychology, 1914, 25, 325-332.
 5. Journal of Psychology, 1914, 25, 325-332.

Foster's objection on the basis of physiological improbability might have been well-based twenty-five years ago; however, present-day physiology does recognize a certain inertia and spontaneity in the nervous system. Forbes, in his discussion of "The Mechanism of Reaction",²⁰ speaks of "after discharge, i.e., persistence of responses after stimulation has ceased, often for several seconds." He further explains the term as "The continuous discharge of impulses in the motor neurons for several seconds after the cessation of afferent stimulation...."

As a matter of fact, the physiological foundation of perseveration has been assumed for some time, especially, in the last decade or so, during which physiological sophistication has grown in psychology. For some investigators, the physiological counterpart of the psychological phenomena forms a necessary part of the definition of the term. It is especially characteristic of a fairly recent tendency to reduce concepts to their "ultimate physical or physiological truths or entities." One of these investigators says it rather clearly and concisely:²¹

Perseveration may be defined as the tendency of a set of neurons, once excited, to persist in the state of excitation autonomously, showing resistance to any change in this state.

Pavlov,²² in referring to the kind of perseveration observed in mental patients, speaks of a "....pathological inertness in the motor area of the cortex."

The entire host of experimenters with the electro-encephalograms must

²⁰ C.Murchison, (ed.) The Foundations of Experimental Psychology, 147-151, 1929.

²¹ H.H.Jasper, "Is Perseveration a Functional Unit Participating in all Behavior Processes," Journal of Social Psychology, 1931, 2, 28-51.

²² I.P.Pavlov, "An Attempt at a Physiological Interpretation of Obsessional Neurosis and Paranoia," Journal of Mental Science, 1934, 80, 190.

Forster's objection on the basis of physiological impossibility might have been well-based twenty-five years ago; however, present-day physiology does recognize a certain inertia and spontaneity in the nervous system. Forster, in his discussion of "The Mechanism of Reaction,"²⁰ speaks of "after discharge, i.e., persistence of response after stimulation has ceased, often for several seconds." He further explains the term as "The continuous discharge of impulses in the motor neurons for several seconds after the cessation of afferent stimulation...."

As a matter of fact, the physiological foundation of perseveration has been assumed for some time, especially in the last decade or so, during which physiological sophistication has grown in psychology. For some investigators, the physiological counterpart of the psychological phenomena forms a necessary part of the definition of the term. It is especially characteristic of a fairly recent tendency to reduce concepts to their "ultimate physical or physiological truths or entities." One of these investigators says it rather clearly and concisely:²¹

Perseveration may be defined as the tendency of a set of neurons, once excited, to persist in the state of excitation autonomously, showing resistance to any change in this state.

²² Pavlov, in referring to the kind of perseveration observed in mental patients, speaks of a ".....pathological hardness in the motor area of the cortex."

The entire host of experiments with the electro-encephalogram must

²⁰ C. Forster, (ed.) The Foundations of Experimental Psychology, 1911, 1939.
²¹ H. H. Jasper, "In Perseveration a Functional Unit Participating in all Behavior Processes," Journal of Social Psychology, 1931, 2, 28-31.
²² I. I. Pavlov, "An attempt at a Physiological Interpretation of Obsessive Neurosis and Paranoia," Journal of Mental Science, 1934, 80, 190.

recognize a certain automaticity and spontaneity of nervous action in the brain, as demonstrated by the action potentials in their sundry variations. The waves (brain waves), as concluded by various investigators, "are due to a spontaneous beat of an area of the occipital cortex concerning with pattern vision; when the cells of this area are not engaged in visual activities they tend spontaneously to discharge impulses at a fixed rate."²³

Moreover, studies on brain potentials do not only definitely establish the spontaneity of central nervous activity, but they also show the perseverative tendency in the cortical activity ("Berger Rhythm") itself. Since light abolishes the rhythm of cortical electrical activity, "Perseveration time is, therefore, the time between the removal of the stimulus (in this case, light) and the reappearance of the potential rhythm."²⁴

Experimenting on the basis of this assumption, the investigators found significant (though as yet uninterpreted) individual differences in perseveration time. Differences in perseveration time by this method were obtained by the same investigators when meaningful and meaningless words were presented to the subjects.²⁵

On the basis of the above presented statements and results from modern physiology and fields of psychological investigation it is safe to conclude that there is in existence a spontaneous activity in the central and, consequently, in the peripheral nervous system. Since that spontaneous activity does not exist directly in the nervous system and since persev-

²³ C.H. Best and N.B. Taylor, The Physiological Basis of Medical Practice, 1404, 1937.

²⁴ L.E. Travis and R.J. Knott, "Brain Potential Studies of Perseveration 1. Perseveration Time to Light," Journal of Psychology, 1936, 3, 97-100.

²⁵ Idem, "Brain Potential Studies of Perseveration 11. Perseveration Time to Visually Presented Words," Journal of Experimental Psychology, 1937, 21, 353.

recognize a certain susceptibility and specificity of nervous action in the brain, as demonstrated by the action potentials in their sensory receptors. The waves (brain waves), as concluded by various investigators, are due to a spontaneous beat of an area of the occipital cortex connected with pattern vision; when the cells of this area are not engaged in visual activities they tend spontaneously to discharge impulses at a fixed rate.²³ Moreover, studies on brain potentials do not only definitely establish

the spontaneity of central nervous activity, but they also show the perseverative tendency in the cortical activity ("after rhythm") itself. Since light abolishes the rhythm of electrical activity, the reaction time is, therefore, the time between the removal of the stimulus (in this case, light) and the reappearance of the potential rhythm.²⁴ Experimentation on the basis of this assumption, the investigators

found significant (though as yet unexpressed) individual differences in perseveration time. Differences in perseveration time by this method were obtained by the same investigators when meaningful and meaningless words were presented to the subject.²⁵

On the basis of the above presented statements and results from modern physiology and theory of psychological investigation it is safe to conclude that there is in existence a special and activity in the central and, consequently, in the peripheral nervous system. Since that spontaneous activity does not exist directly in the nervous system and since perse-

²³ C. E. Hart and H. E. Taylor, "The Physiological Basis of Mental

Exercises," 1904, 1927.

²⁴ H. E. Hart and H. E. Taylor, "Brain Potential Studies of Perse-

vation I. Perseveration Time to Light," Journal of Psychology, 1935, 20, 100.

²⁵ Ibid., "Brain Potential Studies of Perseveration II. Perseveration

Time to Visually Presented Words," Journal of Experimental Psychology, 1937,

21, 208.

eration is a directly observable characteristic even of that spontaneous activity itself, it appears to be quite clear that psychological phenomena grouped under the term perseveration rather gain their validity in terms and in the light of modern physiology.

D. Perseveration in Psychiatry

The phenomenon of perseveration has been noticed as a constant characteristic of some mental disorders long ago. It was not named as such until Neisser (See page 4) introduced the term. Even after the introduction of the term, Kraeplin as well as other psychiatrists grouped perseveration as part of the clinical picture in schizophrenia.²⁷

Wir finden bei der schizophrenie auch manchman eine besondere Art der Perseveration, die allerdings in ihren spezifischen Eigentuemlichkeiten noch nicht recht zu beschreiben ist, obwohl wir annehmen muessen, dass sie sich von anderen Formen unterseheidet. Zufaelliche psychische Vorgaenge, Gedanken in allen Varianten Zusammenhangen immer wieder gedacht und ausgesprochen, 'zu Tode gehetzt.'.....*

He further discusses perseveration in connection with organic psychoses. There the tendency to distinguish it from stereotypy is present; the distinction, however, does not seem too clear.²⁸

Jedefals ist etwas ganz anderes als die Stereotypie die Perseveration der Organischen, die wir am typischsten bei groben

²⁶ E. Kraeplin, (R.M.Barclay, Trans.) Dementia Praecox and Paraphrenia, 21-63, 1919.

²⁷ E. Bleuler, Lehrbuch der Psychiatrie, 61, 1918.

²⁸ Ibid, 110.

*We find in schizophrenia a special kind of perseveration which cannot be yet described, in all its specific characteristics, though we have to assume that it is differentiated from all other kinds. Casual psychic processes, thoughts in all kinds of connections are always thought again and repeated continuously. (Translation---my own.)

Hirnherden, namentlich Aphasikern, sehen. Die Kranken kommen von einem Wort, dass sie eben gehoert oder namentlich gesprochen, nicht ab und wiederholen es, wenn sie etwas anderes sagen wollen, bestaendig. Auch wenn die Patienten eine einfach Handlung ausfuehren, ja bloss etwas denken wollen, geht ihren unter Unstaenden der Impuls gegen ihren Willen in die Bahn der vorhergehenden Handlung oder des unmittelbar vorher gedachten.*

Many additional clinical descriptions of perseveration in psychotics of different and varying diagnoses could be quoted; space will not permit that. But, a clearer line of demarcation between stereotypy and perseveration was drawn at a later date.²⁹

....over activity may consist of a repetition of movements with little or no diversity....Such an oft-repeated movement is known as stereotypy....Perseveration, or the repetition of a recent movement in spite of the patient's effort to produce a new movement, is to be distinguished from stereotypy. It is in its particular instances transient, requires a stimulus, and is associated particularly with cases showing changes in the brain substance.

Thus it appears that perseveration as such is more fluctuating and less constant than the phenomena classified under the heading of stereotypy.

The tendency to persevereate was also observed in mental patients during their subjection to the association test. Thus Bleuler³⁰ and Jung³¹ give detailed descriptions of the tendency to persevereate. Some words are used in reacting continuously to different stimulus words prescribed by the experimenter. However, Jung emphasizes the special connection of the perseverated words with emotional content. Thus, here again, perseveration

²⁹ D.K.Henderson and R.D.Gillespie, A Textbook of Psychiatry, 3rd ed., 83, 1933.

³⁰ Bleuler, op.cit., p.56.

³¹ C. Jung, Collected Papers on Analytical Psychology, 1922.

*Quite different from stereotypy is perseveration which we especially see in cases of aphasia. The patients do not turn away from a word which they recently heard; they repeat it continuously when they even want to say something else. Also, when the patient performs some activity, even if he only wants to think of something, the impulse, under certain circumstances, goes against his desire in the path of the previous activity or of that thought of directly before. (Translation---my own.)

Wiederholte, manchmalige Wiederholungen, die in einem Wort, das ein Element der Wiederholungen enthält, wiederholt werden, sind ein Zeichen für eine Wiederholungsstörung. Diese Wiederholungen sind ein Zeichen für eine Wiederholungsstörung, die in einem Wort, das ein Element der Wiederholungen enthält, wiederholt werden, sind ein Zeichen für eine Wiederholungsstörung.

Many additional clinical descriptions of perseveration in psychotics

of different and varying diagnoses could be quoted; space will not permit

that. But, a clearer line of demarcation between atypical and persever-

ation was drawn at a later date.²²

...over actively may consist of a repetition of movements with little or no alteration. ...which an off-pitched movement is known as stereotypy. ...stereotypy, or the repetition of a movement in the same manner, is a sign of a patient's effort to maintain a new movement, as to be distinguished from stereotypy. It is in the persistent phase of persistent, repeated movements, and is associated particularly with cases showing changes in the brain structure.

Thus it appears that perseveration as such is more frequent and

less constant than the phenomena classified under the heading of stereotypy.

The tendency to perseverate was also observed in mental patients

during their subjection to the association test. Thus Kline²³ and Jung²⁴

give detailed descriptions of the tendency to perseverate. Some words are

used in reacting continuously to different stimulus words presented by the

experimenter. However, Jung emphasizes the special connection of the

perseverated words with emotional content. Thus, here again, perseveration

²² J. H. Robertson and R. F. Gillies, A Textbook of Psychiatry, 1932.

²³ Kline, op. cit., 1931.

²⁴ Jung, Collected Papers on Analytical Psychology, 1932.

*This differs from stereotypy in perseveration which we have seen in cases of aphasia. The patients do not turn away from a word when they repeatedly hear it; they repeat it continuously and even when they are something else. Also, when the patient hears a word, which is not only words to him of something, the patient, who repeats himself, only repeats his desire in the path of the previous activity or at that thought of activity before. (Translators' note.)

is described as a characteristic of a particular state of mental order
and a general mental characteristic.

It is often pointed out that the characteristic of mental order
is the opposite of perseveration, and is often found
with the initial perseveration, and that the two are
independent of each other.

Generally speaking, the characteristic of mental order is
found in the initial perseveration, and is often found
with the initial perseveration, and that the two are
independent of each other.

CHAPTER II

GENERAL EXPERIMENTATION ON PERSEVERATION

It is often pointed out that the characteristic of mental order
is the opposite of perseveration, and is often found
with the initial perseveration, and that the two are
independent of each other.

It is often pointed out that the characteristic of mental order
is the opposite of perseveration, and is often found
with the initial perseveration, and that the two are
independent of each other.

¹ 1915, III.

² S.E. Cassen, Defective Experimental Psychology, 71, 1922.

CHAPTER II

GENERAL EXPERIMENTATION ON
IMITATION OF PRESERVATION

is described as a characteristic of a peculiar state of mentality rather than a general human characteristic.³²

As it often happens that an association of strong feeling tone produces in the experiment a perseveration, with a result that not only the critical association, but also two or three successive associations are imperfectly reproduced.

Generally speaking, the various clinical observations of psychiatrists concerning perseveration adopt empirically the molar view in their descriptions. They see the process as a repetition of an entire pattern of activities rather than a tendency towards perseveration in the simplest and most elementary biophysical processes (molecular view). In the experimental work done by psychiatrists and psychologists the more elementary processes related to perseveration are investigated (See next Chapter). However, the definitive difference between the psychiatric view and understanding of the term and that of the experimental psychological use is shown by Cameron, who says that "Something different from perseveration in the usual psychiatric sense is meant, whether the difference is merely of degree, is yet unclear."³³

One can only state, in this connection, that it is the tacit assumption of the psychologist (attempts to substantiate it have been made) that perseveration is a general characteristic of all human beings, inherent in the nervous system, which, due to organic, known and unknown causes, is more salient and exaggerated in psychotic patients.

³² Ibid, 111.

³³ D.E.Cameron, Objective Experimental Psychiatry, 71, 1935.

is described as a characteristic of a particular state of mind rather than a general characteristic.

As it often happens that an association of ideas follows from a process in the experiment, with a result that not only the original association, but also two or three successive associations are frequently reproduced.

Generally speaking, the various clinical observations of psych-

iatric conditions have been made about experimentally the same view in their descriptions. They are the process as a repetition of an entire pattern of activities rather than a tendency towards preservation in the simplest and most elementary biological processes (biological view). In the experimental work done by psychiatrists and psychologists the more elementary processes related to preservation are investigated (See next Chapter). However, the definite difference between the psychiatric view and understanding of the term and that of the experimental psychological one is shown by Gellert who says that "something different from preservation in the usual psychiatric sense is meant, namely the difference is merely of degree,

in yet another.

One can only state, in this connection, that it is the fact

assumption of the psychologist (attempts to substantiate it have been made) that preservation is a general characteristic of all human beings, inherent in the nervous system, which, due to organic, known and unknown causes, is more salient and exaggerated in psychotic patients.

A. The Dutch School

The attention of a number of investigators was directed, at the beginning of the present century, to the hypothesis advanced by Gross concerning the "Sekundaerfunction."¹ The desire arose to test those interesting observations by means of experimentation under controlled conditions. The first investigator to deal with the secondary function experimentally was Wiersma.² The task before him was quite difficult; he had to invent tests and experimental procedures which would adequately test Gross' theoretical notions and empirical observations. In the main, Wiersma used three tests with which he expected to establish the rightful place of the secondary function in the psychology of individual differences. The tests were as follows: adaptation to light, adaptation to electrical stimulation and color mixture. By means of these methods Wiersma proceeded to investigate the secondary function in manic, normal, melancholic and paranoiac individuals. The underlying rationale of the tests was that length of time for adaptation to light and the strength of the electric stimulation as well as the number of the rotations of the color wheel until grey is seen, are measures of the "Sekundaerfunction." The longer the adaptation time, the greater is the strength of current, and, the fewer the number of revolutions, the greater is the secondary function. The phenomenon of adaptation being a measure of the after effect or lag of the sensory stimulation; it is a sort of sensory "Nachfunction" or inertia of activity, while the electrical adaptation and color disc demonstrate a similar type of inertia. The results of this investigation were in

¹ O. Gross, Die Zerebrale Function, 1902.

² E. Wiersma, "Die Sekuendaerfunktion bei Psychosen," Journal f. Psychol u. Neurologie, 1906, VIII, 1-24.

The attention of a number of investigators was directed, at the beginning of the present century, to the hypothesis advanced by Gross concerning the "sensory function". The desire arose to test those interesting observations by means of experimentation under controlled conditions. The first investigator to deal with the secondary function experimentally was Wierzbicka. The task before him was quite difficult; he had to invent tests and experimental procedures which would adequately test those theoretical notions and empirical observations. In the main, Wierzbicka used three tests with which he expected to establish the right place of the secondary function in the psychology of individual differences. The tests were as follows: adaptation to light, adaptation to electrical stimulation and color mixture. By means of these methods Wierzbicka proceeded to investigate the secondary function in manic, normal, melancholic and epileptic individuals. The underlying rationale of the tests was that length of time for adaptation to light and the strength of the electric stimulation as well as the number of the rotations of the color wheel until gray is seen, are measures of the "sensory function". The longer the adaptation time, the greater is the strength of current, and the fewer the number of revolutions, the greater is the secondary function. The phenomenon of adaptation being a measure of the after effect or lag of the sensory stimulation, it is a sort of sensory "inertia" or lack of activity, while the electrical adaptation and color disc demonstration is another type of inertia. The results of this investigation were in

accord with Gross' views;³ their detailed analysis, however, has to be postponed until the next chapter. Wiersma did not, only, create methods and produce results, but also hinted at new methods and manners of attack upon the problem. In the end of the above-mentioned investigation he also points to some motor phenomena akin to the secondary function, which later became the starting point for the so-called motor perseveration tests. He observed that ".....es schwer ist, eine Bewegung, die man laengeren Zeit in derselben Weise ausgeuebt hat, ploetzlich zu aentern."*⁴ This difficulty in the sudden change of movement is suggested to be due to the secondary function or after effect of the original activity. However, it was not until later that this problem was attacked with motor as well as sensory tests.

Two other Dutch investigators, Heymans and Brugmans⁵ have extended Wiersma's investigations of Gross' theories. The following tests were applied by the authors:

1. Color mixing
2. Limen for flicker
3. Light adaptation
4. Sound adaptation (after loud noise)
5. Motor perseveration
6. Errors in sentences.

The first four tests are obviously sensory and are based on a rationale identical with the one quoted above. The last two tests are motor in nature. In the fifth test, the subject wrote forty letters in the regular manner, then twenty letters in the opposite direction; the time for the last five letters of the forty and the first five of the twenty is

³ Ibid., 18.

⁴ Ibid., 24.

⁵ C. Heymans and H. Brugmans, "Intelligenzpruefungen mit Studierenden," Zeitsch. f. Psych., 1913, 7, 317-331.

* ".....it is difficult to change a movement which one has for some time preferred in the same manner." (Translation--my own.)

accord with Gross' view,⁵ their detailed analysis, however, has to be postponed until the next chapter. Wernke did not only create methods and produce results, but also hinted at new methods and manners of attack upon the problem. In the end of the above-mentioned investigation he also pointed to some motor phenomena akin to the secondary function, which later became the starting point for the so-called motor perseveration tests. He observed that "...as sooner lat, eine Bewegung, die man laengeren Zeit in derselben Weise ausfuehrt hat, desto leichter es kommt." This difficulty in the sudden change of movement is suggested to be due to the secondary function or after effect of the original activity. However, it was not until later that this problem was attacked with motor as well as sensory tests. Two other Dutch investigators, Heymans and Brugmans have extended Wernke's investigations of Gross' theories. The following tests were

applied by the authors:

1. Color mixing
2. Taper for flicker
3. Light adaptation
4. Sound adaptation (after loud noise)
5. Motor perseveration
6. Errors in sentences

The first four tests are obviously sensory and are based on a rationale identical with the one quoted above. The last two tests are motor in nature. In the fifth test, the subject wrote forty letters in the regular manner, then twenty letters in the opposite direction; the time for the last five letters of the forty and the first five of the twenty is

3 min., 15.
4 min., 20.

5. Heymans and Brugmans, "Versuchsergebnisse mit Versuchsgegenstand", Psychologische Zeitschrift, 1912, 7, 214-221.

* "...it is difficult to change a movement which one has for some time performed in the same manner." (Translation--Wernke.)

compared and serves as an index of perseveration or secondary function, since the greater length of time in the second five letters is due, presumably, to the interference of the after function or its after effects (i.e. perseveration).

The sixth test consists of sentences containing similar words in which the persistence of errors in reading the new sentence with similar but not identical words, is measured. Heymans and Brugmans appear to confirm the results of Wiersma and the nature of the secondary function proposed by Gross.⁶ Their statement runs as follows:

Man kann demnach sagen, dass die Ergebnisse saemtlicher vorliegender Tests sehr entschieden, auf eine Bestaetigung, was Otto Gross als Sekundaerfunction bezeichnet hat, und der Nachwirkung von Empfindungen hinweisen.

The experimenters received an average intercorrelation of $\pm .26$. Of course the number of subjects was quite small (fifteen). Spearman⁷ rightfully doubts the significance of the results.

The Dutch School began a series of investigations into the secondary function of the perseveration tendency, devised a number of tests which served as a basis for later investigations and directed their work along two main paths of inquiry in an attempt to answer the following questions:

1. What are the individual differences in regard to the secondary function among normals and psychotics?

2. Do the tests, applied in the measurement of that secondary function, have a common factor; i.e., are they sufficiently the same to

⁶ ibid., 325.

⁷ C. Spearman, The Abilities of Man, 295., 1927.

** One may therefore say that the results of the battery of tests quite decisively confirm what Otto Gross called Secondary function and show the after-effects of sensations. (Translation--my own.)

compared and serves as an index of perseveration or secondary function, since the greater length of time in the second list is due, presumably, to the interference of the first list or its after effects (1933, p. 100).

The sixth test consists of sentences containing similar words in which the persistence of errors in repeating the new sentence with similar but not identical words, is measured. Sayings and sayings appear to conform the results of the test and the nature of the secondary function proposed by Gross. Their statement runs as follows:

Man kann manchmal sagen, dass die Eigenschaften eines Wortes, die es in einem Satz hat, auch in einem anderen Satz sein können, und dass man dann die Eigenschaften des Wortes in einem anderen Satz wiederfindet. (Gross, 1933, p. 100)

The experimenters received an average intercorrelation of .38. Of course, the number of subjects was quite small (fifteen) and it is difficult to judge the significance of the results.

The Dutch School began a series of investigations into the secondary function of the perseveration tendency, devised a number of tests which served as a basis for later investigations and directed their work along two main paths of inquiry in an attempt to answer the following questions:

1. What are the individual differences in regard to the secondary function among normals and psychotics?
2. To the tests, applied in the measurement of this secondary function, have a common factor, i.e., are they sufficiently the same to

G. Gross, 1933, p. 100.
The Psychology of Man, 1933, p. 100.

One may therefore say that the results on the battery of tests give a fairly good picture of the secondary function and show the after-effects of perseveration. (Translation--Gross, 1933, p. 100)

measure the same thing which may run through the various activities?

The results were inadequate and inconclusive; but their stimulating value for future research was great.

B. The English School

Lankes⁸ was the first English pioneer to attack, experimentally, the problem of perseveration. The tests used in his study, were the following: (a) natural rate of tapping; (b) letter writing; (c) cancellation; (d) color disc; (e) drawings; (f) narratives; (g) associative reactions and (h) essays. His results show an intercorrelation, though not a very high one. These, as well as the investigations of Jones⁹ and Bernstein¹⁰ appear to serve the main basis for Spearman's formulation of the "law of inertia" (quoted in Chapter I) and for his faith in the functional unity of perseveration. When more closely observed, the criticism of the conclusions seems to be justified. Shevach offers the following criticism:¹¹

.....The tests were administered as group tests, an inadequate procedure in view of the subtle differentiation involved. In the case of color discs the distance from the exposure slit as well as the angle of vision, was different for different members of the group...

This partial exposition of flaws in experimental procedure and technique due to group testing of perseveration (especially Lankes) raises considerable doubt as to the value of those experimental results.

Jones' battery of tests consisted of the following:¹²

- (1) Writing an S, first repeatedly in the usual way, and then as

⁸ W. Lankes, "Perseveration," British Journal of Psychology, 1915, 7, 387-419.

⁹ C. Spearman, The Abilities of man, 1927.

¹⁰ B. Shevach, "Studies in Perseveration VI. Methods for the Study of Sensory Perseveration," The Journal of Psychology, 3, 381-402.

¹¹ Ibid., 381

¹² C. Spearman, The Abilities of Man, 269, 1927.

measure the same thing which was through the various activities.
The results were inadequate and inconclusive; but their stimulating
value for future research was great.

5. The English School

Lewin⁸ was the first English pioneer to attack, experimentally, the
problem of perseveration. The tests used in his study, were the following:
(a) natural rate of tapping; (b) letter writing; (c) cancellation; (d) color
disc; (e) drawing; (f) narratives; (g) associative reactions and (h) essays.
His results show an intercorrelation, though not a very high one. These,
as well as the investigations of Jones⁹ and Bernstein¹⁰ appear to serve the
main basis for Spearman's formulation of the "law of inertia" (quoted in
Chapter I) and for his faith in the functional unity of perseveration.
When more closely observed, the criticism of the conclusions seems to be
justified. However, the following criticism:¹¹

.....the tests were administered as group tests, an inadequate
procedure in view of the subtle differentiation involved. In the
case of color discs the distance from the exposure slit as well as
the angle of vision, was different for different members of the group...

This partial exposition of flaws in experimental procedure and technique
due to group testing of perseveration (especially Lewin's) raises consider-
able doubt as to the value of these experimental results.

Jones' battery of tests consisted of the following:¹²
(1) Writing on 2. First repeatedly in the usual way, and then as

⁸ Lewin, "Perseveration," British Journal of Psychology, 1925,
17, 257-269.
⁹ C. Spearman, The Abilities of Man, 1927.
¹⁰ A. Bernstein, "Studies in perseveration VI. Methods for the study
of sensory perseveration," The Journal of Psychology, 5, 361-403.
¹¹ Journal of Psychology, 1927, 1927.
¹² Jones, The Psychology of Man, 1927.

it would appear in a mirror.

(2) Writing digits, first in the usual way, and then making the stroke backwards.

(3) Mirror drawing.

(4) First copying prose in the usual way, and then doing it without dotting the i's and crossing the t's. ("it" test). The average positive intercorrelation of these tests employed by Jones was .492, as compared with very low correlations in Lankes' work ("None of the average intercorrelations between the tests exceed plus .29").¹³ The reason for it may be found in the greater homogeneity of Jones' tests involving motor activity.

Bernstein used the following tests in his investigation of 130 children:¹⁴ tapping, S test, backward digits, reverse order letters, the "it" test, re-writing six letters mirror-wise, triangles (first, apex upward and then downward, then alternating), reversal of capital and small letters in copying a passage, copying simple geometrical figures replacing vertical lines by horizontal lines and vice versa; finally, in copying a passage, an a was to be written after every e.

The correlations seem to be lowest in Bernstein's tables. However, when the pool of the tests was correlated with estimates of the subjects' possession or lack of possession of perseverative tendencies, it gave a positive coefficient of .51. Thus, there seemed to be ample validation by an outside criterion. Spearman in his application of the tetrad difference

¹³ B. Shevach, "Studies in Perseveration VI. Methods for the Study of Sensory Perseveration," The Journal of Psychology, ¹⁹³⁵ 3, 381-402.

¹⁴ G. Bernstein, "Quickness and Intelligence," British Journal of Psychology, Mon. Suppl. 1924, VII.

it would appear in a mirror.

(2) Writing digits, first in the usual way, and then making the

stroke backwards.

(3) Mirror drawing.

(4) First copying prose in the usual way, and then doing it without

getting the 'a' and crossing the 't's' (1st test). The average positive
intercorrelation of these tests employed by Jones was .495, as compared
with very low correlations in Langers' work (None of the average inter-
correlations between the tests exceed plus .33%).¹² The reason for it
may be found in the greater homogeneity of Jones' tests involving motor
activity.

Hernstein used the following tests in his investigation of LBO
children:¹³ copying, 2 test, backward digits, reverse order letters, the
1st test, re-writing six letter-mirror-wise, triangles (first, apex upward
and then downward, then alternating), reversal of capital and small letters
in copying a passage, copying simple geometrical figures, tracing vertical
lines or horizontal lines and vice versa; finally, in copying a passage,
and was to be written after every 5.

The correlations seem to be lowest in Hernstein's tables. However,
when the pool of the tests was correlated with estimates of the subjects'
possession of lack of possession of perceptive tendencies, it gave a
positive coefficient of .61. Thus, there seemed to be ample validation by
an outside criterion. Reported in his application of the trend method

12. L. L. Jones, "Studies in Intercorrelation of Tests for the Study
of Language Development," The Journal of Psychology, 1931-32,
14, 341-357.
13. G. Hernstein, "Children and Intelligence," British Journal
of Psychology, vol. 30, 1939, vii.

equation¹⁵ to all of the above-mentioned corrections concludes that ".... in general the whole of the correlations derive from only one factor."¹⁶ This is further substantiated by the insignificant correlations with "g" and with "speed" and "fluency" as demonstrated by Hargraves.¹⁷

Subsequent investigations of perseveration have spread over a wide field. It was the English School, especially, that pursued the following main paths of research and experimentation:¹⁸

(a) The validity of perseveration and development of pertinent psychometric procedures.

(b) The relationship between perseveration and other well known major variables.

(c) The role of perseveration in personality differences.

There seems to be considerable overlap between aspects (b) and (c), also the latter aspect of research has to be assumed as including pathological categories.

However, the rest of this section will largely be devoted to aspect (a) and somewhat to (b), while the entire next chapter is devoted to the abnormal aspect of (c) since it is in line with the subject of the present investigation.

Few renovations have been made in the psychometric technique during the past twenty years. Jones,¹⁹ besides using some of the usual sensory (visual adaptation and color disc) and motor (reverse stroke and "it") tests

¹⁵ C. Spearman, The Abilities of Man, 73-74, 1927.

¹⁶ Ibid, 303.

¹⁷ H.L.Hargreaves, "The Faculty of Imagination," British Journal of Psychology, Mon. Suppl., 1927, 10, 1-74.

¹⁸ K.H.Rogers, "Perseveration," Journal of Mental Science, 1935, 61, 138-144.

¹⁹ L.W.Jones, "An Investigation into the Significance of Perseveration," Journal of Mental Science, 1928, 74, 653-659.

... to all of the above-mentioned correlations...
 in general the whole of the correlation factor...
 This is further substantiated by the following correlation with the
 and the "age" and "illness" as demonstrated by Hargrave.

Subsequent investigations of persistence have shown even a wide
 range. It was the English School, especially, that pursued the following
 main paths of research and experimentation:

- (a) The validity of persistence and development of persistence
- (b) The relationship between persistence and other well known
- (c) The role of persistence in psychological adjustment.

There seems to be considerable overlap between (a) and (b), and
 the latter aspect of research has to be treated as including psychological
 adjustment.

However, the rest of this section will largely be devoted to a report
 of the research in (a), with the entire last section is devoted to the
 research aspect of (b) which is in line with the rest of the section.

The research has been made in the following manner: (a) The
 first twenty years. Jones, 1910, has been the main source of the material
 (which is available and some other) and most (which is available and some other).

10. E. Hargrave, The Psychology of Persistence, 1910, pp. 1-100.
 11. E. Hargrave, The Psychology of Persistence, 1910, pp. 1-100.
 12. E. Hargrave, The Psychology of Persistence, 1910, pp. 1-100.
 13. E. Hargrave, The Psychology of Persistence, 1910, pp. 1-100.
 14. E. Hargrave, The Psychology of Persistence, 1910, pp. 1-100.

also used ideational tests that were somewhat new in character (c.f. Lankes). He used the "noun test" where the subject was to build a number of nouns beginning with a certain letter and then the letter was changed and the observations of the persistence of the previous letter in the new procedure were made.²⁰ The animal test was largely a measure of speed as to how many names of animals can be stated in a given period of time. It is doubtful whether the latter is really a measure of perseveration. The "Blot test" measuring recurring responses does seem to fit into the picture of the definitions of perseveration. Moreover, later investigators on the Rorschach²⁰ seem to confirm the blot value, especially in the case of epileptics. Jones' main objective is not to establish the validity and reliability of his measures, since he gives no intercorrelations, but to correlate the tests with psychiatric categories.

Subsequent work by Stephenson²¹ in which five of the above-mentioned motor tests were used, points out that they have an "average intercorrelation of .40 and satisfy the tetrad criterion for two factors."²²

Application of the same tests was common in the next few years. Epilepsy,²³ the "Resemblance of Twins"²⁴ with regard to perseveration as measured by the motor tests, and the high correlation between siblings²⁵ have all been investigated. Despite the criticism launched against the

²⁰ A. Guirdham, "Rorschach Test in Epileptics," Journal of Mental Science, 1935, 81.

²¹ W. Stephenson, "Studies in Experimental Psychiatry II Some Contacts of p. factor with Psychiatry," Journal of Mental Science, 1932, 78, 315-30.

²² Ibid., 319.

²³ J.H.Ewen, "Perseveration in the Insane Epileptic," Journal of Mental Science, 1930, 76, 537-40.

²⁴ E.P.Yule, "The Resemblance of Twins with Regard to Perseveration," Journal of Mental Science, 1935, 81, 489-501.

²⁵ R.B.Cattell, "Perseveration and Personality--Some Experiments and a Hypothesis," Journal of Mental Science, 1935, 81, 151-67.

also used identical tests that were somewhat new in character (e.g. ladders). He used the "motor test" where the subject was to build a number of houses beginning with a certain letter and then the letter was changed and the observations of the persistence of the previous letter in the new procedure were made. The animal test was largely a measure of speed as to how many names of animals can be stated in a given period of time. It is doubtful whether the latter is really a measure of perseveration. The "slot test" measuring repeating responses does seem to fit into the picture of the definition of perseveration. Moreover, later investigators on the Horschach²⁰ seem to confirm the slot value, especially in the case of epileptics. Jones' main objective is not to establish the validity and reliability of his measures, since he gives no intercorrelation, but to correlate the tests with psychiatric categories.

Subsequent work by Stephenson²¹ in which five of the above-mentioned motor tests were used, points out that they have an "average intercorrelation of .40 and satisfy the tetrad criterion for two factors."²² Application of the same tests was common in the next few years. Epilepsy,²³ the "Resemblance of Twins,"²⁴ with regard to perseveration as measured by the motor tests, and the high correlation between siblings²⁵ have all been investigated. Despite the criticism launched against the

- 20 A. Goldhamer, "Horschach Test in Epileptics," Journal of Mental Science, 1935, 81.
- 21 W. Stephenson, "Studies in Experimental Psychology II Some Correlations of a Factor with Psychiatric," Journal of Mental Science, 1935, 78, 315-30.
- 22 Ibid., 81.
- 23 J.R. Ewen, "Perseveration in the Insane Epileptic," Journal of Mental Science, 1930, 76, 437-40.
- 24 R.F. Yule, "The Resemblance of Twins with Regard to Perseveration," Journal of Mental Science, 1925, 61, 433-36.
- 25 R.B. Cattell, "Perseveration and Personality--Some Experiments and a Hypothesis," Journal of Mental Science, 1935, 81, 151-67.

customary tests²⁶ in which the variation in the size of the letters may upset and destroy the value of the perseveration score, which is based on the ratio of the number of letters produced in the two or more different types of activity, some investigators consider the problem of motor and sensory tests in perseveration settled. Thus, the following statement reveals the confidence of Cattell:²⁷

The validity and reliability of 'p' tests is now clearly on the increase. Nor is there any further reason to doubt that they are measuring a definite group factor 'p' in motor and sensory tests.

It may be noticed that the letter "p" is being used; it is not merely an abbreviation for the term perseveration. With the increase of research on factor analysis, the tendency on the part of previous investigators of "perseveration" is to use "p" instead. This is not merely a tacit and indirect expression of doubt as to whether the tests actually measure perseveration. It means that our present knowledge and definition of perseveration are not quite satisfactory. But, since by the means of the correlation technique, a common factor is found in many tests that claim to measure perseveration, this factor should more adequately receive a less a priori name than perseveration. Stephenson's²⁸ doubts are quite obvious though there is definitely "....a 'p' factor, but its explanation might lie in clearness variation and its laws...."²⁹

This criticism is far from being pessimistic about the tests and about their ability to measure perseveration; it merely shows that the trend

²⁶ D.E.Cameron and T.G.B.Caunt, "Studies in Perseveration," Journal of Mental Science, 1933, 79, 735-745.

²⁷ R.B.Cattell, "Perseveration and Personality--Some Experiments and a Hypothesis," Journal of Mental Science, 1935, 81, 151-67.

²⁸ W.Stephenson, "Perseveration and Character," Character and Personality, 1935, 4, 44-52.

²⁹ Idem, "Studies in Perseveration," Nature, 1934, 134, 860.

constantly tests²⁵ in which the variation in the size of the letters may be-
 not and destroy the value of the perseveration score, which is based on the
 ratio of the number of letters produced in the two or more different types
 of activity, some investigators consider the problem of motor and sensory
 tests in perseveration settled. Thus, the following statement reveals the

confidence of Gestalt:

The validity and reliability of 'p' tests is now clearly on the
 increase. Now is there any further reason to doubt that they are
 measuring a definite group factor 'p' in motor and sensory tests.

It may be noticed that the letter "p" is being used; it is not

merely an abbreviation for the term perseveration. With the increase of

research on factor analysis, the tendency on the part of previous

investigators of "perseveration" is to use "p" instead. This is not merely

a habit and indirect expression of doubt as to whether the tests actually

measure perseveration. It means that our present knowledge and definition

of perseveration are not quite satisfactory. But, since by the means of the

correlation technique a common factor is found in many tests that claim to

measure perseveration, this factor should more adequately receive a less

a priori name than perseveration. Stephenson's²⁶ doubts are quite obvious

though there is definitely ".... a 'p' factor, but its explanation might lie

in cleanness variation and its laws...."²⁷

This criticism is far from being pessimistic about the tests and

about their ability to measure perseveration; it merely shows that the trend

²⁵ D. E. Campbell and T. O. B. Grant, "Studies in Perseveration," *Journal of Mental Science*, 1933, 73, 735-743.

²⁶ W. A. Gestalt, "Perseveration and Personality--Some Experiments and Hypotheses," *Journal of Mental Science*, 1933, 73, 151-67.

²⁷ W. A. Stephenson, "Perseveration and Character," *Character and Personality*, 1933, 4, 44-52.

²⁸ Ibid., "Studies in Perseveration," *Nature*, 1934, 134, 660.

is away from the psychologically defined terminology towards that of "factors" where "w" replaces will and volition, "p"---secondary function and perseveration, et cetera. It is not, of course, merely a change of names. It implies that the term is getting at something more fundamental, biologically and experimentally defined.

C. Experimental Work in the United States

Much less attention was paid to the concept of perseveration in this country. There was, of course, clinical recognition of the phenomenon. Thus, White³⁰ and other psychiatrists and workers in the field of abnormal psychology understood it in the same sense that Neisser³¹ indicated originally. A few experiments, however, follow the English tradition in experimenting with perseveration.

Cushing's study on "A Perseverative Tendency in Pre-School Children"³² approaches and attacks the problem by giving more "natural" tests taken from the children's life-like situations. On the basis of her work, the investigator concludes in favor of a perseverative tendency as follows:³³ "The results seemed to indicate the presence of a common factor pervading the situation plus certain small group factors...."

Of course, it may be a point of contention as to whether the "persistence" of "dropping marbles through a hole into a box" is comparable to the after-sensation and after effects of an activity characteristic of the tests in the British investigations. However, according to the summary-

³⁰ W.A.White, Outlines of Psychiatry, 13th ed., 1932.

³¹ See Chapter I, Section B. present study.

³² H.M.Cushing, "A Perseverative Tendency in Pre-School Children," Archives of Psychology, 1929, 108.

³³ Ibid., 52.

is away from the psychobiologically defined terminology towards that of "factor", where "factors" with and without, "p"---secondary function and preservation, of nature. It is not, of course, merely a change of name. It implies that the term is setting at something more fundamental, biologically and experimentally defined.

3. Experimental work in the United States

More than attention was paid to the concept of preservation in this country. There has, of course, clinical recognition of the phenomenon. Thus, White³⁰ and other psychologists and workers in the field of abnormal psychology understood it in the past years that what was indicated originally. A few experiments, however, follow the English tradition in experimenting with preservation.

Washburn's study on "A perseverative tendency in pre-school children" approaches and attacks the problem by giving more "natural" tests taken from the children's life-like situation. On the basis of her work, the investigation concludes in favor of a perseverative tendency as follows:
"The results seemed to indicate the presence of a common factor pervading the situation plus certain small group factors...."

Of course, it may be a point of contention as to whether the "perseverance" or "drop-out" method through a hole into a box is comparable to the after-association and after effects of an activity characteristic of the tests in the British investigation. However, according to the summary-

30 E. A. White, Journal of Psychology, 1934, 21, 193.
31 See Chapter I, Section A, present study.
32 E. A. Washburn, "A perseverative tendency in pre-school children," Archives of Psychology, 1933, 103.
33 ibid., 1933, 103.

definition of perseveration to be given later, it will appear that Cushing's work is covered by the blanket term of perseveration or perseverative tendency.

The most extensive and more critical study was made by Jasper.³⁴ This investigator employed a long series of tests to his 76 subjects who were students at the University. The tests were sensory, motor and ideational; many of them were of the type employed in the British investigations, while additional reaction-time tests were given. Jasper does not seem to confirm the results of the English experimenters. He seems to get a negligible correlation between the types of tests (ideational, sensory and motor) with regard to perseveration. There is, however, according to him, "....some evidence for the existence of a narrower group factor of motor perseveration....."³⁵ Here, again, the trouble and the objection is not to the term perseveration as such. Its existence and soundness as a psychological entity is granted. But, the adequacy of the present psychometric and experimental methods in the measurements of perseveration is seriously doubted. Also the proof of the consistency of the trait throughout the activities involved in the tests is under attack. Hence, Jasper puts forth the following desiderata:³⁶

Measures must be developed which are sufficiently specific for perseveration. The measures used may be fairly reliable, but not valid measures of perseveration. Some of the measures used in this study were scored so that other factors could easily have masked a considerable portion of the 'perseverative' tendency.

Related to the last portion of Jasper's statement is also one made by

³⁴ H.H. Jasper, "Is Perseveration a Functional Unit Participating in all Behavior Processes," Journal of Social Psychology, 1931, 2, 28-51.

³⁵ Ibid., 48.

³⁶ Ibid., 45.

definition of perseveration to be given later, it will appear that Jasper's work is covered by the blanket term of perseveration or perseverative tendency.

The most extensive and more critical study was made by Jasper.³⁴ This investigator employed a long series of tests to his 78 subjects who were students at the University. The tests were sensory, motor and ideational; many of them were of the type employed in the British investigations, while additional reaction-time tests were given. Jasper does not seem to confirm the results of the English experiments. He seems to get a negligible correlation between the types of tests (ideational, sensory and motor) with regard to perseveration. There is, however, according to him ".....some evidence for the existence of a narrower group factor of motor perseveration....."³⁵ Here, again, the trouble and the objection is not to the term perseveration as such, its existence and soundness as a psychological entity is granted, but the adequacy of the present psychometric and experimental methods in the measurement of perseveration is seriously doubted. Also the proof of the consistency of the trait throughout the activities involved in the tests is under attack. Hence, Jasper cuts forth the following desiderata:³⁶

Measures must be developed which are sufficiently specific for perseveration. The measures used may be fairly reliable, but not valid measures of perseveration. Some of the measures used in this study were scored so that other factors could easily have masked a considerable portion of the 'perseverative' tendency.

Related to the last portion of Jasper's statement is also one made by

³⁴ H. H. Jasper, "Is Perseveration a Unimodal Unit Factor?" in "All Behavior Problems," Journal of Social Psychology, 1931, 3, 22-31.
³⁵ 1931, 3, 22-31.
³⁶ 1931, 3, 22-31.

Dorcus.³⁷ In the conclusions of his study, he says that "...such a general factor based upon tissue condition or neural tendency may be obscured by desires and volition."³⁸ This last objection is not quite so sound since "volition and desires" may be related to that neural tendency or be a part of it. Further discussion of this point will take us too far afield--- into the deep waters of metaphysics.

The most recent comprehensive series of studies on perseveration comes from the Harvard Psychological Clinic. Kendig and Shevach³⁹ have initiated that series of investigations, the former investigator--on conative processes involved in perseveration, while the second investigator (Shevach) outlined the aim of his study as follows:⁴⁰

(a) To determine whether sensory perseveration is a general characteristic of personality, or whether it varies in degree from one sensory experience to another.

(b) To determine the constancy of perseveration under similar and dissimilar conditions.

(c) To determine the influence of age upon the phenomenon.

Shevach employed a great many sensory tests (eleven). Some of them had to be discarded as a result of the inadequacy of laboratory conditions. The number of small experimental groups gave a rather low average inter-correlation of the tests used. However, some points of importance have been established in the following conclusions:⁴¹

1. Sensory perseveration of a given sensory experience manifests marked individual differences.

2. Sensory perseveration is manifested as a general characteristic

³⁷ R.M.Dorcus, "Is there a 'Unit-Trait' of Volitional perseveration?" Journal of General Psychology, 1935, 13, 345-356.

³⁸ Ibid., 354.

³⁹ I.Kendig and B.J.Shevach, "Studies in Perseveration, I. A Study Survey of Researches in Perseveration," The Journal of psychology, 1936, 3, 223-230.

⁴⁰ ibid., 228.

⁴¹ B.J.Shevach, "Studies in Perseveration, VII. Experimental Results of Tests for Sensory Perseveration," The Journal of Psych., 1933, 403-427.

37 In the discussion of his study, he says that "...such a general

factor based upon status condition or mental tendency may be observed by
38 desired and volition." This last objection is not quite so sound since

volition and desire may be related to that mental tendency or be a part
of it. Further discussion of this point will take us too far afield--

into the deep waters of metaphysics.

The most recent comprehensive series of studies on perseveration

39 comes from the Harvard Psychological Clinic. Reading and Eysenck have

indicated that series of investigations, the former investigator--on cognitive
processes involved in perseveration, while the second investigator, Eysenck

40 outlined the aim of his study as follows:

- (a) To determine whether sensory perseveration is a general
characteristic of personality, or whether it varies in degree from one
sensory experience to another.
- (b) To determine the constancy of perseveration under similar and
dissimilar conditions.
- (c) To determine the influence of age upon the phenomenon.

Eysenck employed a great many sensory tests (eleven). Some of them
had to be discarded as a result of the inadequacy of laboratory conditions.

The number of small experimental groups gave a rather low average inter-
correlation of the tests used. However, some points of importance have been

41 established in the following conclusions:

1. Sensory perseveration of a given sensory experience manifests
marked individual differences.
2. Sensory perseveration is manifested as a general characteristic

42 M. Eysenck, "Is there a Unit-Test of Volitional Perseveration?"
Journal of General Psychology, 1955, 12, 345-352.

43 I. A. Eysenck, "Studies in Perseveration, I. A Study
of Research in Perseveration," *The Journal of Psychology*, 1955, 40, 353-362.

44 I. A. Eysenck, "Studies in Perseveration, VII. Experimental Results
of Tests for Sensory Perseveration," *The Journal of Psychology*, 1955, 40, 363-372.

in some individuals and varies in degree from one sensory experience to another amongst other individuals.

3. Psychoneurotic tendencies constitute the direct factor which underlies the variability in the degree of perseveration.

4. The degree of perseveration for a given sensory experience is constant from time to time.

Thus, the reliability and manifestations of individual differences on the basis of the sensory tests is quite definite; also, the relationship between variability and psychoneurotic tendencies and their effect upon the establishment of sensory perseveration as a general characteristic is to be noted. Another conclusion included in the final summary of the study should prove of great interest. He says:⁴²

Extreme perseveration and extreme non-perseveration predispose an individual to psychic invalidism.

The status of the individuals in regard to psychoneurotic tendencies was established by means of various personality tests (Thurstone, Woodworth and Bernreuter.) The variability was finally interpreted as a result of a basically low perseverative tendency which is "disrupted" and varies from situation to situation, while the stable persons, who are somewhat high in the middle range of perseveration scores, give a more consistent picture all around. The variations in the former depend upon the emotional value of the situation.

These last results do not appear very clear-cut. To some extent, the last conclusions seem to confirm the results obtained earlier by Rogers⁴³ that "High and low 'Perseverators' show these tendencies ('disparities' between intelligence and school achievement) to a greater degree than did those children who come within the median ranges of the 'Perseveration'

⁴² ibid., 426.

⁴³ K.H. Rogers, "A Relationship Between Perseveration and School Achievement," Psychological Bulletin, 1930, 30, 163-69.

in some individuals and varies in degree from one sensory experience to another among other individuals.
 3. Psychomotoric tendencies constitute the direct factor which explains the variability in the degree of perseveration.
 4. The degree of perseveration for a given sensory experience is constant from time to time.

Thus, the reliability and manifestations of individual differences on the basis of the sensory tests is quite definite, also, the relationship between variability and psychomotoric tendencies and their effect upon the establishment of sensory perseveration as a general characteristic is to be noted. Another conclusion included in the final summary of the study should prove of great interest. He says:

Extreme perseveration and extreme non-perseveration predominate in individual to psychic instability.

The status of the individuals in regard to psychomotoric tendencies was established by means of various personality tests (Thurstone, Woodworth and Barnet). The variability was finally interpreted as a result of a basically low perseverative tendency which is "disrupted" and varies from situation to situation, while the stable persons, who are somewhat high in the middle range of perseveration scores, give a more consistent picture all around. The variations in the former depend upon the emotional value of the situation.

These last results do not appear very clear-cut. To some extent, the last conclusions seem to confirm the results obtained earlier by Rogers that high and low perseverators show these tendencies ('disruptive' between intelligence and school achievement) to a greater degree than did these children who come within the median range of the 'perseveration'.

distribution." The confirmation, of course, depends upon the agreement with the assumption that "discrepancies" are indicators of instability and psychoneurotic tendencies.

Some more work done in the psychiatric field in this country will be reviewed in greater detail in a more appropriate place. It seems, however, that, on the whole, investigators in the United States neglected the problem when compared with the output of their colleagues in England. The work is more critical. On the one hand, a narrow group factor of motor perseveration seems to be warranted by the evidence;⁴⁴ while on the other hand, there seems to be (with certain qualifications) a group factor of sensory perseveration which is a fairly stable and reliable measure.⁴⁵ Some relation between perseveration and neurotic tendencies has been established. But, little or no work was done with the motor and sensory tests and their measurements of perseveration in psychotic subjects, since the above experimenters employed normal subjects only.

D. Summary of Experimental Methods

All the methods (psychometric) employed in the measurement of perseveration or secondary function may be classified under the following headings:

1. Motor tests
2. Sensory tests
3. Ideational tests

under the first class come different kinds of motor tests, tests which involve or regard switching from one activity to another similar to it, break-

⁴⁴ Jasper, op.cit., 2, 28-51.

⁴⁵ Shevach, op.cit., 3, 403-427.

disturbance." The correlation, of course, depends upon the agreement with the assumption that "disturbances" are indicators of instability and psychotic tendencies.

Some more work done in the psychiatric field in this country will be reviewed in greater detail in a more appropriate place. It seems, however, that, on the whole, investigators in the United States neglected the problem when compared with the output of their colleagues in England. The work is more critical. On the one hand, a narrow group factor of motor perseveration seems to be warranted by the evidence;⁴⁴ while on the other hand, there seems to be (with certain qualifications) a group factor of sensory perseveration which is a fairly stable and reliable measure.⁴⁵ Some relation between perseveration and neurotic tendencies has been established, but, little or no work was done with the motor and sensory tests and their measurements of perseveration in psychotic subjects, since the above experiments employed normal subjects only.

Summary of Experimental Methods

All the methods (psychometric) employed in the measurement of perseveration or secondary function may be classified under the following

categories:

1. Motor tests
2. Sensory tests
3. Ideational tests

Under the first class come different kinds of motor tests, tests which involve or regard switching from one activity to another similar to it, break

⁴⁴ Jaber, op. cit., p. 28-31.
⁴⁵ Jaber, op. cit., p. 403-427.

ing away from well-habituated activities, "and, newly instituted mental sets."⁴⁶

The sensory tests include adaptation phenomena, threshold sensitivities and reaction times.

The ideational tests contain "fluency" tests (animals, ink-blots, et cetera), memory (recall) tests, tests dealing with spontaneous recurrence of mental elements of association---tests, essays, et cetera) and interference tests, when the person's mental set is shown to interfere with the one following it (Noun test and others of similar nature.)

As a result of this great expansion in the field of testing perseveration, the intercorrelation of the multitude of tests is in doubt. At least, the studies thus far quoted are not unanimous in this respect. There seems to be, however, some agreement as to the intercorrelation of these tests within each one of the groups (motor, sensory and ideational).

The definition of perseveration, or as it became in psychology---mental inertia, is quite broad and inclusive and seems to include a multitude of phenomena. Both the confusion in the definition of the term as well as the host of experiments in widely differing directions, caused the broadening of its meaning. The list of the following phenomena most adequately summarizes what is meant by perseveration today.⁴⁷

1. Tendency of ideas to spontaneous recurrence
2. Sensory lag or after effect
3. Interference effects of preceding mental processes with present

⁴⁶ W. Stephenson, "An Introduction to So-Called Motor Perseveration Tests," British Journal of Educational Psychology, 1934, IV, 186-207.

⁴⁷ I. Kendig and B.J. Shevach, "Studies in Perseveration 1. A Survey of Researches in Perseveration." The Journal of Psychology 1936, 223-230.

processes.

4. Tendency to continue working at a task when there is no objectively defined end.

Generally speaking, continuance seems to be the cardinal principle involved in perseveration; all the rest of the phenomena are akin, caused by and related to it.

E. Some Theoretical Considerations

We have examined here, in some detail, the theoretical and experimental foundations of a problem. It appears, as a result of this examination, that perseveration is a rather elusive trait which is not entirely independent of other variables. But, "strict independence is neither a vital nor even a reasonable requirement for human traits."⁴⁸ Some independence of the trait, some group-factor is evident. The evidence came from two directions: first,---from theoretical definitions which defined tests, which in turn, measured perseveration; an a priori method. Second,---from the experimental data and employment of statistical analysis to the discovery of general factors---the a posteriori method.

Now, it seems relatively unimportant, for all practical purposes, whether we use the term perseveration and argue for its being a universal trait occurring in all our activities, or whether we say that there is a "p" factor which was statistically derived from a great variety of tests which had their genesis and original stimulation in the theory of perseveration. They both measure the same phenomena; both approach each other from different sides of the road.

⁴⁸ G.W.Allport, Personality, A Psychological Interpretation, p.430, 1937.

4. Tendency to continue working at a task when there is no

objectively defined end.

Generally speaking, constancy seems to be the cardinal principle involved in perseveration; all the rest of the phenomena are either caused by and related to it.

5. Some Theoretical Considerations

We have examined here, in some detail, the theoretical and experimental foundations of a problem. It appears, as a result of this examination, that perseveration is a rather elusive trait which is not entirely independent of other variables. But, strict independence is neither a vital nor even a reasonable requirement for human traits.

Good independence of the trait, some group-factor is evident. The evidence came from two directions: first, from theoretical definitions which defined tests, which in turn, measured perseveration; and a priori

method. Second, from the experimental data and employment of statistical analysis to the discovery of general factors--the statistical method. Now, it seems relatively unimportant, for all practical purposes,

whether we use the term perseveration and argue for its being a universal trait occurring in all our activities, or whether we say that there is a factor which was statistically derived from a great variety of tests which had their genesis and original stimulation in the theory of

perseveration. They both measure the same phenomenon, both approach each other from different sides of the road.

Thus, the existence of the phenomena here described is undeniable; the tests used are fallible and a greater amount of "p" in new tests is to be desired,---greater, but not complete independence of other factors is to be sought. The relation to individual differences, character and personality types, is further to be investigated and explored (See Chapter III).

CHAPTER III

EXPERIMENTATION WITH PSYCHOTIC MATERIAL

Thus, the existence of the phenomena here described is undoubted;
the facts noted are reliable and a greater amount of "p" in new tests is to
be expected,--greater, but not complete independence of other factors in
the result. The relation to individual differences, character and
personality types, is further to be investigated and explored (See Chapter

1. Personality Correlated with the Various Diagnoses [Psychiatric]

Most of the investigations, whose work was discussed in the previous chapter, have been interested in establishing some truth concerning the validity of the various diagnoses. It is necessary to realize that this correlation of personality with the various diagnoses was not a simple matter. It is necessary to realize that the correlation of personality with the various diagnoses was not a simple matter. It is necessary to realize that the correlation of personality with the various diagnoses was not a simple matter.

CHAPTER III

EXPERIMENTATION WITH PSYCHOTIC MATERIAL

The investigation in this chapter is concerned with the correlation of personality with the various diagnoses. It is necessary to realize that the correlation of personality with the various diagnoses was not a simple matter. It is necessary to realize that the correlation of personality with the various diagnoses was not a simple matter. It is necessary to realize that the correlation of personality with the various diagnoses was not a simple matter.

The table on page 33 gives a composite picture of the results. The results are not quite identical with those reported by Sperry, who reported that the number of subjects who were in the various diagnoses was not a simple matter. It is necessary to realize that the correlation of personality with the various diagnoses was not a simple matter.

1. S. Gross, The Psychology of the Individual, 1933.
2. See Chapter II, Section 1, for a more detailed study.
3. S. Gross, "The Psychology of the Individual," Journal of Psychology, 1933, VIII, 1-24.

CHAPTER III

EXPERIMENTATION WITH PSYCHIC MATERIAL

A. Perseveration Correlated with the Various Diagnoses (Psychiatric)

Most of the investigators, whose work was discussed in the previous chapter, have been interested in establishing some truth concerning the unitary function of perseveration, its reliability and validity. A by-product of those investigations was the correlation of perseveration with various character and personality traits, especially along the introversion-extroversion continuum. A number of experiments, however, was devoted to the correlation and investigation of various diagnostic-psychiatric groups in regard to perseveration. It must be recalled that, after all, the genesis of the term is in psychiatry and that Gross'¹ early hypothesis definitely linked it to the greater secondary function in melancholia and the lesser in mania. Upon the basis of results obtained in investigations of abnormal subjects, generalizations were made and extended to the field of normal typology. The number of researches dealing with psychotic subjects does not seem to present a very clear picture as to the linkage of perseveration with the several diagnoses used in the classification of mental patients. Thus Wiersma², using three sensory tests, seems to have confirmed the theoretical predictions uttered by Gross.

The table on page 33 gives us a composite picture of Wiersma's³ results. The results were not quite adequately presented by Spearman, who reported that the number of subjects employed in Wiersma's investigation consisted of nine manics, eleven normals and eighteen melancholics. It appears that this was the largest number in any single experiment (color

¹ O. Gross, Die Zerebrale Sekundaerfunktion, 1902.

² See Chapter II, Section A, present study.

³ E. Wiersma, "Die Sekundaerfunktion bei Psychosen," Journal f. Psychol u. Neurologie, 1906, VIII, 1-24.

A. Perseveration Correlated with the Various Diseases (Psychiatric)

Most of the investigators, whose work was discussed in the previous chapter, have been interested in establishing some truth concerning the history function of perseveration, its reliability and validity. A by-product of those investigations was the correlation of perseveration with various character and personality traits, especially along the introversion continuum. A number of experiments, however, was devoted to the correlation and investigation of various diagnostic-psychiatric groups in regard to perseveration. It must be recalled that, after all, the genesis of the term is in psychiatry and that Gross,¹ early hypothesized definitely linked it to the greater secondary function in melancholia and the lesser in mania. Upon the basis of results obtained in investigations of abnormal subjects, generalizations were made and extended to the field of normal typology. The number of researchers dealing with psychotic subjects does not seem to present a very clear picture as to the linkage of perseveration with the several diagnoses used in the classification of mental patients. Thus Wierma,² using three sensory tests seems to have confirmed the theoretical predictions uttered by Gross.

The table on page 33 gives us a composite picture of Wierma's results. The results were not quite adequately presented by Spearman, who reported that the number of subjects employed in Wierma's investigation consisted of nine manics, eleven normals and sixteen melancholics. It appears that this was the largest number in any single experiment (color

1 G. Gross, Die Verweilungs-Erscheinungen, 1902.
2 See Chapter II, Section A, present study.
3 E. Wierma, "Die Sekundärfunktion bei Psychosen," Journal f. Psychol. u. Neurol., 1903, VIII, 1-24.

TABLE I

SENSORY PERSEVERATION IN NORMALS AND PSYCHOTICS (Wiersma)			
Experiment	Manic	Normal	Melancholic
Electric Adaptation	0 3	39.6 5	205.5 4
Color Disc	27.2 .11	15.7 9	12.2 18
Light Adaptation	33.5 7	102.9 4	233.2 18

TABLE II

JONES' RESULTS

TEST	MANIA	NORMALS	MELANCHOLIA
Adaptation I	54	41	38
Adaptation II	45	31	25
Red Color Green	443	662	541
Blue Yellow	447	674	463

TABLE I

PRIMARY INVESTIGATION OF HORMONES AND HYPERACTIVITY (Wistar)

Experiment	Mean	Normal	Abnormal
Adaptation	0	23.6	200.8
Color	27.7	14.7	12.2
Light	22.8	102.8	223.2

TABLE II

LOWESS RESULTS

TEST	MEAN	LOWESS	ADAPTATION
Adaptation I	24	21	26
Adaptation II	25	21	23
Color	242	262	261
Light	242	272	232

disc), but not in the other two. The number of subjects upon whose results the means of the table are based is given in the corner of each square and is evidently much smaller for any single tests than those reported by Spearman. On the basis of these results, the manic - normal - melancholic order of increasing perseveration was established; thus, supposedly confirming Gross' contentions. Two points of criticism, however, must be leveled at the conclusiveness of these results. First, the number of subjects employed is really, statistically, inadequate and hence, the conclusiveness of the results is impaired. Second, from the present writer's own analysis of Wiersma's data and from what is to be expected when comparing manics with melancholics, the observation must be made that there is a considerable age difference between the two groups. The manics as a group were much younger than the melancholics; most of the former were in their twenties, while the melancholic group was composed of much older persons.

These critical notes are to be born in mind since their significance for the present investigation is of great importance.

About twenty years after Wiersma's study was published a similar investigation with apparently contradictory results was undertaken and completed.⁴ The old sequence of manic-- normal - melancholic in regard to the degree of perseveration is altogether reversed on the basis of two of the sensory tests (Adaptation and Color Disc). The table on page 33 presents Jones' composite results on the two tests.⁵

Thus, manics, according to Jones' results, need a longer adaptation

⁴ C. Spearman, The Abilities of Man, 294, 1927.

⁵ L.W.Jones, "An Investigation into the Significance of Perseveration," Journal of Mental Science, 1926, 74, 653-659.

direct, but not in the other two. The number of subjects upon whose results
 the means of the table are based is given in the corner of each square and
 is evidently much smaller for any single test than those reported by
 Greenman. On the basis of these results, the manic - normal - melancholic
 order of increasing perseveration was established; thus supposedly con-
 firming Greenman's conclusions. Two points of criticism, however, must be
 leveled at the conclusiveness of these results. First, the number of
 subjects employed is really, statistically, inadequate and hence, the
 conclusiveness of the results is weakened. Second, from the present
 writer's own analysis of Greenman's data and from what is to be expected when
 comparing manics with melancholics, the perseveration must be made that there
 is a considerable age difference between the two groups. The manics as a
 group were much younger than the melancholics; most of the former were in
 their twenties, while the melancholic group was composed of much older
 persons.

These critical notes are to be born in mind since their significance
 for the present investigation is of great importance.
 About twenty years after Warram's study was published a similar
 investigation with apparently contradictory results was undertaken and
 completed.⁴ The old sequence of manic - normal - melancholic in regard to
 the degree of perseveration is altogether reversed on the basis of two of
 the sensory tests (Adaptation and Color Disc). The table on page 22
 presents Jones' composite results on the two tests.⁵
 Thus, manics, according to Jones' results, need a longer adaptation

⁴ E. Greenman, *The Abilities of Man*, 1937.
⁵ L.W. Jones, "An Investigation into the Significance of Perseveration,"
Journal of Mental Science, 1938, 74, 658-669.

time than do melancholics and normals, hence, they are greater perseverators (since the after-effect persists longer). Also, the manics lost the two colors of the sectors of the color disc much sooner, after fewer revolutions, than did the melancholics; in other words the manic patients needed fewer rotations of the disc per minute before the individual colors blended, and, consequently, the persistence of the sensations of the individual colors was longer and more effective. Thus, the manics, again, were the greater perseverators. The sequence in regard to sensory perseveration seems to be reversed, i.e., melancholia - normals - mania. The rest of the study (Jones) seems to be independent of the first part. Motor and ideational tests are used to measure "fluency" and it is concluded that "....the results bear out the general view as to the "fluency" of associations in mania and the lack of associations in melancholia."⁶ The question of the relation between fluency and perseveration seems to be quite unsettled, since the author dares to conclude tentatively that "In any case these preliminary results seem to weaken the validity of the view that manic patients are non-perseverators in sensory or motor tests."⁷ Another point has to be emphasized in this connection: the intercorrelation of the tests was quite high in melancholia (motor and ideational) but very low in mania. Whether "large variations in type" may account for it is not yet known.

This study, then, contradicts the results of Wiersma on the basis of sensory tests only. It presents a very unclear picture regarding the existence of the same factor in motor and ideational tests--neither have differences in degree of perseveration between the several psychiatric

⁶ Ibid., 657.

⁷ Ibid., 659.

time than do melancholics and normals. Hence, they are greater perseverators (since the after-effect persists longer). Also, the manics lost the two colors of the sectors of the color disc much sooner, after fewer revolutions, than did the melancholics; in other words the manic patients needed fewer rotations of the disc per minute before the individual colors blended, and, consequently, the persistence of the individual colors was longer and more effective. Thus, the manics, again,

were the greater perseverators. The asynergy in regard to sensory perseveration seems to be reversed, i.e., melancholia - normal - mania. The rest of the study (Jones) seems to be independent of the first part. Motor and ideational tests are used to measure "flimsey" and it is concluded that "....the results bear out the general view as to the 'flimsey' of associations in mania and the lack of associations in melancholia."

The question of the relation between flimsey and perseveration seems to be quite unsettled, since the author seems to conclude tentatively that "in any case these preliminary results seem to weaken the validity of the view that manic patients are non-perseverators in sensory or motor tests."

Another point has to be emphasized in this connection: the intercorrelation of the tests was quite high in melancholia (motor and ideational), but very low in mania. Whether "large variations in type" may account for it is not yet known.

This study, then, contradicts the results of Wiersma on the basis of sensory tests only. It presents a very unclear picture regarding the existence of the same factor in motor and ideational tests - rather than differences in degree of perseveration between the several psychomotor

categories, been clearly demonstrated. Mania and melancholia have been separated on the basis of "fluency", but the connection and relation between fluency and perseveration is definitely not indicated, and proof of its existence is wanting.

A more extensive attack upon the problem, according to the advice of Spearman,⁸ was undertaken by Stephenson⁹ some years later. His experiments were strictly confined to the motor type of test which was used throughout the series of investigations with psychotic and normal subjects.

The table on page 37 is a composite of the results obtained in the above-mentioned study.¹⁰ About 150 subjects were used; among them were some normals and a great many psychiatric categories and sub-classes. According to the median score on "p", for every group (see table) the manic patients are lowest, next are the paranoiacs and other delusionals, epileptics, normals, depressive manic-depressives, melancholics and dementia praecox with high "p".

We have not mentioned one group included in the table since it needs special explanation. The "dementia supervening cases" with a comparatively high "p" score are those cases which show great deterioration (probably due to brain lesions) and which include some cases of general paresis and other cases described in the next paragraph:¹¹

The results that we have to show agree to some extent, in that manic cases involving supervening dementia do not give low 'p' scores; we find,

⁸ C. Spearman, "The Tenth Maudsley Lecture," Journal of Mental Science, 1929, 75, 357-370.

⁹ W. Stephenson, "Studies in Experimental Psychiatry 11 Some Contacts of p. - factor with Psychiatry," Journal of Mental Science, 1932, 78, 315-330.

¹⁰ Ibid.

¹¹ Ibid., 325.

unaffected, have clearly demonstrated. Manic and melancholic states have
regarded on the basis of "illness", but the connection and relation between
illness and personality is definitely not indicated, and proof of the
connection is wanting.

A more extensive attack upon the problem, according to the advice
of Spemann,⁸ was undertaken by Stephenson⁹ some years later. His
experiments were strictly confined to the motor type of test which was used
throughout the series of investigations with psychotic and normal subjects.
The table on page 37 is a composite of the results obtained in the
above-mentioned study.¹⁰ About 100 subjects were used; among them were
some normals and a great many psychiatric categories and sub-classes.
According to the median score on "p", for every group (see table) the manic
patients are lowest, next are the paranoiacs and other delusional, schizo-
phrenics, normals, depressive manic-depressives, melancholics and obses-
sionists with high "p".

We have not mentioned one group included in the table since it needs
special explanation. The "depressive subparanoiacs" with a comparatively
high "p" score are those cases which show great deterioration (probably
due to brain lesions) and which include some cases of general paresis and
other cases described in the next paragraph.¹¹

The results that we have to show are, in that order,
the cases involving subparanoiacs, manic-depressives, normals, schizo-

⁸ C. Spemann, "The Psychomotoric Test," *Journal of Experimental Psychology*, 1923, 25, 337-370.
⁹ W. Stephenson, "A Study in Experimental Psychology," *Journal of Experimental Psychology*, 1923, 25, 315-350.
¹⁰ Table 1, page 37.
¹¹ Table 2, page 38.

TABLE III

P-SCORES FOR PSYCHOTICS AND NORMALS (Stephenson)

SUBJECTS	NUMBER	MEDIAN P SCORES	SCORE AND ESTIMATE
Paranoiacs	21	2.5	9
Other delusionals		4	
Dementia Praecox with high P		17	45
Other Cases of Dementia Praecox	25	4	9
Melancholics	21	9	30
Manic-depressive Psychosis (Depressive Phase)	14	7	23
Maniacs	9	1	5
Epileptics	14	5	14
Dementia Supervening Cases	19	12	32
Normal	20	5	16

TABLE VII

I-SCORES FOR PSYCHOTICS AND NORMALS (continued)

SUBJECT	NUMBER	MEDIAN I-SCORE	SCORE AND ESTIMATE
Paranoid	21	2.5	3
Other Schizophrenia		4	
Manic-depressive with manic phase		17	42
Other cases of Manic-depressive	25	4	3
Manic-depressive	21	9	20
Manic-depressive (Depressive Phase)	14	7	22
Manic-depressive	9	1	5
Psychotics	14	5	14
Manic-depressive Superior group	13	12	28
Normal	20	6	18

however, that other manics, of good 'g' ability generally do obtain very low 'p' scores, and, in our estimation, have clearly non-'p' characteristics.

Thus we actually have two types of manics---the non-deteriorated and the deteriorated which cannot be classed with the former. Moreover, according to Stephenson's optimistic views, the correspondence between a high p-score and degree of deterioration in manics is so definite and consistent that "....the use of the 'p' tests in the manic group....might be of service in objectively helping to decide upon matters of mental reduction and deterioration."¹² Hence, the statement concerning the manic as non-perseverators is robbed of its universality and must be qualified.

The dementia praecox group also presents very interesting results. The distribution of the perseveration scores of this group would produce practically a bi-modal curve. There is one group (see Table II, page 33) with a very high "p" score and another group with an extremely low "p" score ---nearly approaching the delusionals and paranoiacs. Hence, this lack of uniformity and point to point correlation with the diagnoses may seemingly cause us considerable doubt concerning the value of the "p" tests in this connection. Stephenson foresaw this and investigated the problem more thoroughly in his next study.¹³ This study confirmed the clinical observations made in the earlier study---namely: that high "p" correlates with low accessibility of the patient and vice versa, low "p" with greater accessibility. Thus, in cases of dementia praecox the state of psychosis can fairly well be determined on the basis of the "p" tests. Stephenson's own statement concerning this matter goes as follows:¹⁴

¹² Ibid., 326.

¹³ W. Stephenson, "Studies in Experimental Psychiatry: III. p.-score and Inhibition for High p.-Praecox Cases," Journal of Mental Science, 1932, 78, 908-920.

¹⁴ Ibid., 913.

however, that other members of good 'e' ability generally to obtain very low 'e' scores, and, in our estimation, have clearly non-'e' characteristics.

Thus we actually have two types of manic-depressed and the deteriorated which cannot be classed with the former. Moreover, according to Stephenson's optimistic views, the correspondence between a high p-score and degree of deterioration in manic is no definite and consistent that the use of the 'p' tests in the manic group.... might be of service in objectively helping to decide upon matters of mental retention and deterioration. Hence, the statement concerning the manic as non-perservers is robbed of its universality and must be qualified.

The dementia praecox group also presents very interesting results. The distribution of the perseveration scores of this group would produce practically a bi-modal curve. There is one group (see Table II, page 35) with a very high 'p' score and another group with an extremely low 'p' score. Hence, this lack of uniformity and point to point correlation with the diagnoses may seemingly cause us considerable doubt concerning the value of the 'p' tests in this connection. Stephenson foresaw this and investigated the problem more thoroughly in his next study. This study confirmed the clinical observations made in the earlier study--namely: that high 'p' correlates with low accessibility of the patient and vice versa, low 'p' with greater accessibility. Thus, in cases of dementia praecox the state of psychosis can fairly well be determined on the basis of the 'p' tests. Stephenson's own statement concerning this matter goes as follows:

14 Ibid., 315.
15 Ibid., 320.
16 Ibid., 320.
17 Ibid., 320.
18 Ibid., 320.
19 Ibid., 320.
20 Ibid., 320.
21 Ibid., 320.
22 Ibid., 320.
23 Ibid., 320.
24 Ibid., 320.
25 Ibid., 320.
26 Ibid., 320.
27 Ibid., 320.
28 Ibid., 320.
29 Ibid., 320.
30 Ibid., 320.
31 Ibid., 320.
32 Ibid., 320.
33 Ibid., 320.
34 Ibid., 320.
35 Ibid., 320.
36 Ibid., 320.
37 Ibid., 320.
38 Ibid., 320.
39 Ibid., 320.
40 Ibid., 320.
41 Ibid., 320.
42 Ibid., 320.
43 Ibid., 320.
44 Ibid., 320.
45 Ibid., 320.
46 Ibid., 320.
47 Ibid., 320.
48 Ibid., 320.
49 Ibid., 320.
50 Ibid., 320.
51 Ibid., 320.
52 Ibid., 320.
53 Ibid., 320.
54 Ibid., 320.
55 Ibid., 320.
56 Ibid., 320.
57 Ibid., 320.
58 Ibid., 320.
59 Ibid., 320.
60 Ibid., 320.
61 Ibid., 320.
62 Ibid., 320.
63 Ibid., 320.
64 Ibid., 320.
65 Ibid., 320.
66 Ibid., 320.
67 Ibid., 320.
68 Ibid., 320.
69 Ibid., 320.
70 Ibid., 320.
71 Ibid., 320.
72 Ibid., 320.
73 Ibid., 320.
74 Ibid., 320.
75 Ibid., 320.
76 Ibid., 320.
77 Ibid., 320.
78 Ibid., 320.
79 Ibid., 320.
80 Ibid., 320.
81 Ibid., 320.
82 Ibid., 320.
83 Ibid., 320.
84 Ibid., 320.
85 Ibid., 320.
86 Ibid., 320.
87 Ibid., 320.
88 Ibid., 320.
89 Ibid., 320.
90 Ibid., 320.
91 Ibid., 320.
92 Ibid., 320.
93 Ibid., 320.
94 Ibid., 320.
95 Ibid., 320.
96 Ibid., 320.
97 Ibid., 320.
98 Ibid., 320.
99 Ibid., 320.
100 Ibid., 320.

....in agreement with the picture of psychiatric change from inaccessibility and very uncontrolled behavior to greater accessibility and quickness, there is shown marked decrease in 'p'-score.

The author even concludes that the "p" tests may serve as a "barometer" for psychiatric condition. Elaborate theories are brought to the fore as a result of the data obtained, however, we shall confine ourselves to the actual results based on experimental work and make our own deductions later in this paper.

The manic-depressives seem to occupy a middle position in regard to the "p" score, while the melancholics appear highest. Thus, it appears that Wiersma's arrangement of manic, normal, melancholic, in regard to perseveration and in contradistinction to Jones' results, still stands. However, an attachment of a number of qualifications (dementia supervening cases) is necessary. Other investigations,¹⁵ carried out around the same time, seem to corroborate Stephenson's and, consequently, Wiersma's view in regard to perseveration in melancholia.

Parenthetically, it may be interesting, also, to observe that the results concerning epileptics in the above quoted studies hardly agree with other studies comparing normals on the same level in regard to perseveration. Ewen¹⁶ obtained results that are diametrically opposed to those and prove distinctly the epileptics to be much higher perseverators than normals. Of course, Ewen only used one motor test while the others were sensory (2) and ideational (2) tests. The results led him to conclude that they show "....that perseveration is a feature of the peculiar mental state

¹⁵ J.W. Pinard, "Their Relation to Certain Psychopathic Conditions and to Introversion," British Journal of Psychology, 1932, 23, 114-127.

¹⁶ J.H. Ewen, "Perseveration in the Insane Epileptic," Journal of Mental Science, 1930, 76, 537-540.

.....in agreement with the picture of psychiatric change from manic-depressive illness and very uncontrolled behavior to greater normality and quietness, there is shown marked decrease in "p"-score.

The author even concludes that the "p" tests may serve as a "barometer" for psychiatric condition. Elaborate theories are brought to the fore as a result of the data obtained, however, we shall confine ourselves to the actual results based on experimental work and make our own deductions later in this paper.

The manic-depressive seem to occupy a middle position in regard to the "p" score, while the melancholics appear highest. Thus, it appears that Witkowsky's arrangement of manic, normal, melancholic, in regard to perseveration and its contribution to Jones' results is still valid. However, an attachment of a number of qualifications (dementia and reversing cases) is necessary. Other investigations¹⁵ carried out around the same time, seem to corroborate Witkowsky's and, consequently, Witkowsky's view in regard to perseveration in melancholia.

Technically, it may be interesting, also, to observe that the results concerning epilepsies in the above quoted studies hardly agree with other studies concerning epilepsy on the same level in regard to perseveration. When¹⁶ obtained results that are diametrically opposed to those and prove distinctly the epilepsies to be much higher perseverators than normals. Of course, when only used one factor test while the others were sensory (2) and intellectual (3) tests. The results led him to conclude that "they were"....that perseveration is a feature of the particular mental state

¹⁵ J. H. Jones, "Witkowsky's Relation to Certain Psychiatric Conditions and to Introversions," British Journal of Psychology, 1932, 25, 114-127.
¹⁶ J. H. Jones, "Perseveration in the Epileptic," Journal of Mental Science, 1930, 76, 227-240.

of insane epileptics between attacks...."¹⁷ One may wonder whether the type of tests in one investigation and the different type in the other may account for the disagreement in results and conclusions.

Throughout the studies, thus far reviewed, there was no mention of age control. The question may arise as to whether variations in age do not mask considerably the actual results. Stephenson confesses this weakness: "...age differences and the like are perhaps focal points of criticism."¹⁸

A later investigation¹⁹ seems to have brought the fact of age differences more to the foreground. Even though the Table²⁰ IV below does not show "striking" differences in "p"-scores due to age differences, some increase with age (general tendency) is evident. That increase

TABLE IV

Average	Number	P - Score
21.2	9	45.9
26.5	14	50.5
32.6	18	50.0
44.1	11	50.6

Cattell's Results

comes to an even level at the age of 26 or so, but, what happens beyond 44.1 is not yet known. There may be a much greater increase in the 50's

¹⁷ Ibid., 540

¹⁸ W. Stephenson, "Studies in Experimental Psychiatry II Some Contacts of p.-factor with Psychiatry," Journal of Mental Science, 1932, 78, 330.

¹⁹ R.B. Cattell, "On the Measurement of 'Perseveration,'" British Journal of Educational Psychology, 1935, 5, 76-92.

²⁰ Ibid., 87.

of inane epigrams between attacks....¹⁷ One may wonder whether the type of tests in one investigation and the different type in the other may account for the disagreement in results and conclusions.

Throughout the studies thus far reviewed, there was no mention of age control. The question may arise as to whether variations in age do not mask considerably the actual results. Stephenson counters this weakness: "...age differences and the like are perhaps focal points of criticism."¹⁸

A later investigation¹⁹ seems to have brought the fact of age differences more to the foreground. Even though the Table IV below does not show "attitude" differences in "a"-scores due to age differences, some increase with age (general tendency) is evident. That increase

TABLE IV

Average	Number	F - Score
31.3	3	48.3
30.3	14	50.3
32.4	18	50.0
44.1	11	50.5

Castro's Results

comes to an even level at the age of 20 or so, but, what happens beyond 22 is not yet known. There may be a much greater increase in the 20's

¹⁷ W. Stephenson, "Attitudes in Experimental Psychology," *Journal of Experimental Psychology*, 1925, 20, 330.
¹⁸ W. Stephenson, "Attitudes in Experimental Psychology," *Journal of Experimental Psychology*, 1925, 20, 330.
¹⁹ W. Stephenson, "Attitudes in Experimental Psychology," *Journal of Experimental Psychology*, 1925, 20, 330.

and 60's just as there is a considerable increase from childhood to adulthood²¹ or from early childhood to puberty.²²

This "focal point of criticism" must be born in mind, since it is also focal and of considerable import in the present study.

A comparatively novel experimental approach to the study of perseveration was recently demonstrated in the work of Mays²³ and Shipley.²⁴ Mays starts out with clinical observation of the great tendency toward perseveration on the part of catatonic patients and makes the following statement:

Since most of the clinically observed perseverations are reactions dependent on a 'voluntary' control, the question may be raised whether such repetitive activity extends likewise to the processes under autonomic control.²⁵

The question was answered in the affirmative by the data obtained in the study in which the psychogalvanic response to certain sound patterns was used as a measurement of perseveration. "Catatonics showed two or more times as much perseveration as normals in all comparisons."²⁶

Shipley's²⁷ results are based on four tests of autonomic perseveration which was measured by means of the psychogalvanometer. The tests were as follows:

1. Resistance to adaptation

²¹ J.W.Pinard, "Tests of Perseveration," British Journal of Psychology, 1932, 23, 5-19.

²² M.Dybowski, "Perseveration as a Measure of Negativism, and its Estimation by Means of Photographs," Kwartolnik Psychologiczny, 1935, VI, 221-236.

²³ L.L.Mays, "Studies in Catatonia V. Perseverational tendencies in Catatonic Patients," The Psychiatric Quarterly, 1934, 8, 728.

²⁴ W.C.Shipley, "Studies of Catatonia VI. Further investigation of the Perseverative Tendency," The Psychiatric Quarterly, 1934, 8, 736-744.

²⁵ Mays, loc.cit., 728.

²⁶ Ibid., 735.

²⁷ Shipley, loc.cit., 742.

and 20% less as there is a considerable difference observed in adult-
hood²¹ or from early childhood to young life.²²

This "local point of existence" must be born in mind, since it is

also local and of considerable import in the present study.

A comparatively novel experimental approach to the study of

perseveration was recently demonstrated in the work of Mays²³ and Shipley.²⁴

Mays started out with clinical observation of the great tendency toward

perseveration on the part of catatonic patients and makes the following

statement:

"Since most of the clinically observed perseverations are reactions
dependent on a 'voluntary' control, the question may be raised whether
such repetitive activity extends likewise to the processes under
autonomic control."²⁵

The question was answered in the affirmative by the data obtained

in the study in which the psychogalvanic response to certain sound patterns

was used as a measurement of perseveration. "Catatonics showed two or

more times as much perseveration as normals in all comparisons."²⁶

Shipley's²⁷ results are based on four tests of autonomic persever-

ation which was measured by means of the psychogalvanometer. The tests

were as follows:

1. Resistance to adaptation

21. W.W. Fernald, "Tests of Perseveration," British Journal of Psychology, 1932, 25, 5-19.
22. W. Dykowitz, "Perseveration as a Measure of Negativity, and its Estimation by means of Photograms," Experimental Psychology, 1933, 26, 321-332.
23. L.L. Mays, "Studies in Catatonia V. Perseverational Tendencies in Catatonic Patients," The Psychiatric Quarterly, 1934, 8, 722.
24. L.L. Shipley, "Studies of Catatonia VI. Further Investigation of the Perseverative Tendency," The Psychiatric Quarterly, 1934, 8, 323-344.
25. Mays, loc.cit., 722.
26. Ibid., 723.
27. Shipley, loc.cit., 722.

2. Conditionability
3. Resistance to experimental extinction
4. Irradiation

The combined "p" scores in the table below seem to establish a sequence which is in agreement with that of Stephenson. The final results of the combined "p" scores are as follows:²⁸

TABLE V.

Schizophrenia	Manic-depressive	Psychoneurosis	Normal
91	80	72	53

Shipley's Results

On the whole, the consistency between the tests is quite good and the schizophrenia - manic-depressive - psychoneurotic - normal sequence may be considered as established. However, neither Shipley nor Stephenson gives measures of reliability which would consolidate their results. A mere comparison between means, especially between small groups, is of little significance. The sampling error may be so great, that in absence of critical ratio values, the results cannot be accepted as truly indicative of definite tendencies.

B. Points of Disagreement

The last section was quite serviceable in exhibiting to us a

²⁸ Ibid., 742.

3. Conditionability

2. Resistance to experimental extinction

1. Irradiation

The combined "p" scores in the table below seem to establish a sequence which is in agreement with that of Stephenson. The final results of the combined "p" scores are as follows:

TABLE V.

Psychoneurotic	Manic-depressive	Schizophrenic	Normal
72	80	91	35

Shipley's Results

On the whole, the consistency between the tests is quite good and the schizophrenic - manic-depressive - psychoneurotic - normal sequence may be considered as established. However, neither Shipley nor Stephenson give measures of reliability which would consolidate their results. A more comparison between means, especially between small groups, is of little significance. The sampling error may be so great, that in absence of critical ratio values, the results cannot be accepted as truly indicative of definite tendencies.

B. Points of Disagreement

The last section was quite extensive in exhibiting us a

considerable number of disagreements between investigators and their experimental results. Those disagreements lead to an inconsistent and a faulty understanding of the respective psychiatric categories by means of perseveration measurements; this inconsistency also complicates and hinders our understanding of perseveration itself. This relationship between psychiatric categories and classifications and degrees of perseveration must be cleared up. What, in brief, are the bones of contention?

1. In the first place, regarding the manic - normal - melancholic sequence, the results of Jones and Wiersma, discussed above, are diametrically opposed to each other. Consequently, the following questions will have to be answered:

(a) Are the melancholics of Wiersma and Jones of the same category?

This may have a partly negative answer; there is some historical evidence that early in this century Wiersma bunched depressives and involuntional melancholics in the same group. Jones, however, must have used only involuntional melancholics in accordance with present usage of the term. Hence, there may be a discrepancy as a result of the different types employed.

(b) Some of the high "p" manic patients of Jones may have had "dementia supervening" described by Stephenson. Thus, what is, actually, the difference between manics and demented manics?

(c) Since there are age differences, among the patients, which were not taken into consideration as a variable, what is the effect of those differences, if any?

2. In the second place, the relationship between the psychoneuroses and perseveration is still an open one. Shevach showed a relationship

considerable number of disagreements between investigators and their
 experimental results. These disagreements lead to an inconsistent and a
 faulty understanding of the respective psychiatric categories of manic
 depression and melancholia; this inconsistency also complicates and hinders
 our understanding of pervasiveness itself. This relationship between
 psychiatric categories and classifications and degrees of pervasiveness
 must be cleared up. What in brief are the points of contention?

1. In the first place, regarding the manic - normal - melancholic
 spectrum, the results of Jones and Kistner, discussed above, are diametrically
 opposed to each other. Consequently, the following questions will have to
 be answered:

(a) Are the relationships of Kistner and Jones of the same category?
 This may have a partly negative answer; there is some historical
 evidence that early in this century Kistner punched depressives and
 involutional melancholics in the same group. Jones, however, must have
 used only involutional melancholics in accordance with present usage of the
 term. Hence, there may be a discrepancy as a result of the different types
 employed.

(b) Some of the high "p" manic patients of Jones may have had
 "depressive superimposed," described by Stephenson. Thus, what is, actually,
 the difference between manics and demented manics?

(c) Since there are age differences among the patients, which were
 not taken into consideration as a variable, what is the effect of these
 differences, if any?

2. In the second place, the relationship between the psychomotor
 and pervasiveness is still an open one. Shvach showed a relationship

between psychoneurotic tendencies and variability in perseveration, while Shipley considered hospitalized psychoneurotics greater perseverators than normals.

In this connection, two points are to be born in mind:

(a) We do not know whether normals with psychoneurotic tendencies are related to stable normals as psychoneurotics are to "normals" in general.

(b) The two investigators used widely differing methods in the measurement of perseveration. Sensory tests were used by one (Shevach), while the psychogalvanic response was used by the other (Shipley).

3. In the third place, the relationship between schizophrenia and manic-depressive psychoses in regard to perseveration is still not clear. Shipley considers the schizophrenics higher perseverators than manic-depressives, while Stephenson presents a different sequence in order of increasing perseveration: manic - "low p" dementia praecox - manic depressive - "high p" praecox. The following questions suggest themselves:

(a) Are not the high "p" and low "p" groups in schizophrenia due to the various types within the general category?

(b) Is there a consistently high perseverative tendency in catatonics when compared with other schizophrenics in accordance with the clinical picture?

(c) How are the various states in the manic-depressive psychoses to be compared with the different classification in schizophrenia?

Answers to all these questions may constitute a program of rather extensive research in the field of experimental psychology and psychiatry. This resumé concerning the "frontiers of ignorance" is the underlying

between psychomotoric tendencies and variability in perseveration, while Shipley considered hospitalized psychomotoric groups as psychomotoric than normals.

In this connection, two points are to be born in mind.
(a) We do not know whether normals with psychomotoric tendencies are related to stable normals as psychomotorics are to "normals" in general.

(b) The two investigation methods used with Shipley's method in the measurement of perseveration. Generally tests were made by one (Shipley), while the psychomotoric response was used by the other (Shipley).
3. In the third place, the relationship between schizophrenia and manic-depressive psychosis in regard to perseveration is still not clear. Shipley considers the schizophrenia high perseverators than manic-depressive, while Stephenson presents a different opinion in order of increasing perseveration: manic - "low" dementia praecox - manic depressive - "high p" processor. The following questions suggest themselves:
(a) Are not the high "p" and low "d" groups in schizophrenia due

to the various types within the general category?
(b) Is there a consistently high perseverative tendency in catatonics when compared with other schizophrenia in accordance with the clinical picture?

(c) How are the various types in the manic-depressive psychosis to be compared with the different classification in schizophrenia?
Answers to all these questions may constitute a program of rather extensive research in the field of experimental psychology and psychiatry.
This research concerning the "function of perseveration" is the underlying

raison d'etre of the present study. It gives the stimulus to the inquiry that is outlined in the next section, which proposes to answer some of the questions that were brought to the fore.

C. Aim of Present Study

It would be an Herculean task to attempt a reply to all the questions raised in the last as well as in the previous sections. In the present study, we shall attempt to answer or "settle" a number of those problems - not all of them. Our interest concentrates, largely, around the dementia praecox and manic-depressive groups, since these are to some extent still terra incognita in the field of psychiatry and some light shed on them would be of great value. Also, according to many authorities the above groups are merely extremes of the normal introversion-extroversion types; since that is the case, our understanding of normals may consequently be increased.

It is, therefore, the aim of this study to investigate the following points:

1. The relationship between age and the present measures of perseveration.
2. A comparison in degree of perseveration between the two psychiatric categories---dementia praecox and manic-depressive psychoses.
3. Comparison, in regard to perseveration, between the sub-groups of dementia praecox (catatonic, hebephrenia, paranoia, et cetera).
4. The degree of perseveration in the two phases (manic and depressive) of the manic depressive psychosis.
5. The relationship between perseveration and introversion -extro-

reasons of the present study. It gives the stimulus to the inquiry that is outlined in the next section, which proposes to answer some of the questions that were brought to the fore.

3. Aim of Present Study

It would be an overstatement to attempt a reply to all the questions raised in the last as well as in the previous sections. In the present study, we shall attempt to answer or "testify" a number of these problems - not all of them. Our interest concentrates, largely, around the domestic pressure and male-feminine groups, since these are to some extent still some lacunae in the field of psychiatry and some light shed on them would be of great value. Also, according to many authorities the above groups are merely extremes of the normal introversion-extraversion types; since that is the case, our understanding of normals may somewhat be increased.

It is, therefore, the aim of this study to investigate the following points:

1. The relationship between age and the present concepts of introversion.
2. A comparison in degree of introversion between the two psychiatric categories -- female and male-feminine groups.
3. Comparison, in regard to introversion, between the two groups of domestic pressure (introverted, dependent, etc. cases).
4. The degree of introversion in the two present female and male-feminine groups.
5. The relationship between introversion and introversion-extraversion.

version.

6. Do the "p" tests measure the same thing?
7. What is the relation between "improvement" and the "p" score?

CHAPTER IV

EXPERIMENTAL METHOD AND REASONING

version.

- 2. To the "Q" tests measure the same thing?
- 3. What is the relation between "improvement" and the "Q" tests?

4. Subjective Data

All the subjects employed in this study were patients drawn from the wards of the Boston State Hospital. The management of this hospital extended to the present investigation its facilities and laboratory facilities. All the subjects were males between the ages of 18 and 30. Many of them were in the wards of the hospital in which the laboratory was located, others were brought by an attendant from other buildings because a "doctor wanted to see them." Some personalities were more open and sociable. Some of the patients are "Tubercular" in any case.

CHAPTER IV

EXPERIMENTAL METHODS AND PROCEDURES

There were two experimental groups. Group A consisted of 22 individuals who were in the wards of the hospital and were of mild-to-depressed depressive states. The preliminary results and final data experimentation with this group (see Chapter V).

The second group (B) consisted of 22 individuals who were in the wards of the hospital and were of mild-to-depressed depressive states. The preliminary results and final data experimentation with this group (see Chapter V).

There was also a small number (7) of people patients in the ward group. These patients were quite normal and were called out of the ward quite well. They were called to come freely to the hospital grounds.

The notes were taken up in the hospital files, with an eye for

VI. RESULTS

EXPERIMENTAL METHOD AND PROCEDURE

A. Subjects Used

All the subjects employed in this study were patients taken from the wards of the Boston State Hospital. The management of this hospital extended to the present investigator its hospitality and laboratory facilities. All the subjects were males between the ages of 19 - 60. Many of them were in the wards of the building in which E's laboratory was located, others were brought by an attendant from other buildings, because a "doctor wanted to see them." Some parole patients came over upon request. None of the patients was "forced" in any way.

There were two experimental groups---A and B. Group A consisted of 32 individuals among whom were 22 cases of dementia praecox and 10 cases of manic-depressive (depressive state). The preliminary results are based upon experimentation with this group. (See Chapter V.)

The second group (B) consisted of 80 individuals among whom were 25 manic-depressives (both phases) and the rest were cases of dementia praecox of a great many varieties---catatonics, paranoics, hebephrenics, simple praecoxes and some whose grouping was undetermined. Among this group there was a number of individuals (10) whose cooperation and reliability of report were unsatisfactory, and, therefore, do not count in the final results.

There was also a small number (7) of parole patients in the above group. These patients were quite well oriented and were taking care of themselves quite well. They were allowed to move freely on the hospital grounds.

The cases were looked up in the Hospital files, with an eye for unanimous diagnosis and elimination of complicating factors like brain

All the subjects employed in this study were patients taken from the wards of the Boston State Hospital. The management of this hospital attended to the present investigation for its hospital and laboratory facilities. All the subjects were males between the ages of 18 - 50. Many of them were in the wards of the building in which E's laboratory was located, others were brought by an attendant from other buildings, because a "doctor wanted to see them." Some parole patients came over upon request. None of the patients was "forced" in any way.

There were two experimental groups---A and B. Group A consisted of 32 individuals among whom were 22 cases of dementia praecox and 10 cases of manic-depressive (depressive state). The preliminary results are based upon experimentation with this group. (See Chapter V.)

The second group (B) consisted of 50 individuals among whom were 22 manic-depressive (both phases) and the rest were cases of dementia praecox of a great many varieties---catatonic, paranoiac, hebephrenic, simple praecox and some whose grouping was undetermined. Among this group there was a number of individuals (10) whose cooperation and reliability of report were unsatisfactory, and, therefore, do not count in the final results.

There was also a small number (7) of parole patients in the above group. These patients were quite well oriented and were taking care of themselves quite well. They were allowed to move freely on the hospital grounds.

The cases were looked up in the hospital files, after an eye for unsatisfactory diagnosis and elimination of complicating factors like brain

injury, arteriosclerosis, symptoms of senility, organic changes, alcoholism and the like. In other words, the desire to try to get "clear-cut" diagnostic groups was dominant in the choice of the subjects. Also, an approximate matching of ages between the praecoxes and depressives was kept in mind.

B. The Preliminary Rapport

The author felt that with the type of subjects used in this study, the establishment of some sort of rapport was an essential part of the experimental technique and procedure. Since the battery of tests was not completely of the "objective" type in which the cooperation and report of the subject was not essential, this rapport was quite necessary. E tried to overcome the following attitudes which are typical of a great many hospitalized psychotic patients:

1. Fear of doctors, E being considered one of them in general; (injections, et cetera.)

2. Fear as part of paranoid, psychotic delusions---"someone is going to put something over on them."

3. Resentment of suspicion that there is something wrong with them. They need no treatment; they "don't want to be touched;" "they are perfectly all right physically and mentally."

4. Fear of instruments---especially electrical.

The attitudes listed above are but a few of the most common ones. Each patient may also have some pet aversions and fears which are none too conducive to regular experimental procedure. In order to overcome the obstacles of the nature described above, it was necessary to introduce a

...ly, anhedonia, symptoms of anxiety, organic changes, nihilism
the like. In other words, the desire to try to get "clean-cut"
...was dominant in the choice of the subjects. Also, an
...of ages between the processes and depressive was
...in mind.

The Preliminary Report

The author felt that with the type of subjects used in this study,
...of some sort of report was an essential part of the
...and procedure. Since the battery of tests was not
...of the "objective" type in which the cooperation and report of
...was not essential, this report was quite necessary. I tried
...the following attitudes which are typical of a great many
...psychotic patients:

1. Fear of doctors, as being considered one of them in general;
...at ceters.)
2. Fear as part of paranoid, psychotic behavior---"someone is
...over on them."
3. Resentment of suspicion that there is something wrong with
...They need no treatment; they "don't want to be troubled;" they are
...physically and mentally."
4. Fear of instruments---especially electrical.

The attitudes listed above are but a few of the most common ones.
Each patient may also have some but variations and fears which are more or
...connected to regular experimental procedure. In order to overcome the
...of the nature described above, it was necessary to introduce a

preliminary conversational period with the patient, preceding the actual experimental procedure and the administration of the tests.

When S is first admitted to the laboratory room, the apparatus cannot be readily seen, (behind screen) and he is invited to sit at the table at which E is seated. The E begins some sort of conversation with S. The conversation does not follow any definite routine, but includes the following items: - length of hospitalization; subject's relatives and their visits; his occupation, or lack of it, on the hospital grounds; educational and economic history; and, if the subject is sufficiently recovered, possibilities of his dismissal from the hospital. If S is talkative, he is permitted to talk and lead to different subjects for a period of fifteen minutes; if he is not talkative, many questions are asked in a casual and none too insistent manner. During this preliminary period E manages (in most cases) to establish some favorable attitudes or overcome negative tendencies, wholly or in part, characteristic of the tested patient.

For some very depressed patients it is necessary to change the mode of response. For instance, instead of expecting a very depressed patient to say, "I feel it now," he is told to take his hands off the chair (electrodes) as soon as he "feels the tingle." Likewise, in the adaptation experiment, after no verbal response is elicited, S is instructed to tap on the table "as soon as he sees the round spot of light." Thus the expected measurement is determined (i.e., adaptation time), though the customary verbal, introspective report is not used. Quite often it was necessary to take numerous measures in order to determine the subject's true adaptation time.

preliminary conversational period with the patient, preceding the actual experimental procedure and the administration of the tests.

When S is first admitted to the laboratory room, the experimenter cannot be readily seen, (behind a screen) and he is invited to sit at the table at which E is seated. The E begins some sort of conversation with S. The conversation does not follow any definite routine, but includes the following items: - length of hospitalization; subject's relatives and their visits; his occupation, or lack of it, on his hospital grounds; educational and vocational history; and, if the subject is sufficiently recovered, hospitalization of his classmates from the hospital. If S is catatonic, he is permitted to talk and lead to different subjects for a period of fifteen minutes. If he is not catatonic, many questions are asked in a casual and more or less insistent manner. During this preliminary period E attempts (in most cases) to establish some favorable attitude or overcome negative tendencies, wholly or in part, characteristic of the catatonic patient.

For some very depressed patients it is necessary to change the mode of response. For instance, instead of expecting a very depressed patient to say, "I feel as now," he is told to take his hands off the chair (introduction) as soon as he feels the tingling. Likewise, in the adaptation experiment, after no verbal response is elicited, S is instructed to tap on the table "as soon as he feels the round spot of light." Thus the expected measurement is determined (i.e., adaptation time), though the customary verbal, introspective report is not used. Quite often it was necessary to take numerous measures in order to determine the subject's true adaptation time.

In the motor tests, tremors, lack of motor coordination, inability to perform delicate movements due to brain injury or other organic reasons, as well as inappropriate comprehension of instructions, hindered in obtaining adequate results in one hundred per cent of the cases. Only the "sure" tests were used in the results; the doubtful material was altogether eliminated. Hence, there is a number of subjects who have satisfactorily completed one or more tests, but not the entire battery; this will be adequately taken into account in the discussion of the results. (See Chapter V).

C. Tests Employed in Study

In order to test the intragroup correlation of the sensory and motor tests and their intergroup correlation with one another, the two traditional types are represented in this study. Two of the most representative motor and two of the sensory tests were chosen. There is no particular justification for the particular choice in each case, except for the fact that from the point of view of the experimenter's experience they would lend themselves to fairly good application with the subjects (psychotics) who were to be used. Other tests were also used (See Section, 3,4) with smaller groups of patients, since excellent cooperation and achievement in all the tests was hardly to be expected.

1. Motor Tests

Both of these tests were used in Stephenson's¹ studies with psychotics. However, there are some modifications here, in the manner of perseveration

¹ W. Stephenson, "Studies in Experimental Psychiatry: III. p.-score and inhibition for High p.-Praecox Cases," Journal of Mental Science, 1932, 78, 908-920.

In the motor tests, tremor, lack of motor coordination, inability to perform delicate movements due to brain injury or other organic reasons, as well as inappropriate comprehension of instructions, hindered in obtaining adequate results in one-handed test of the hands. Only the "pure" tests were used in the present; the doubtful material was either eliminated. Hence, there is a number of subjects who have satisfactorily completed one or more tests, but not the entire battery; this will be adequately taken into account in the discussion of the results. (See Chapter 7).

2. Tests Employed in Study

In order to test the intrapersonal correlation of the sensory and motor tests and their intergroup correlation with one another, the two traditional types are presented in this study. Two of the most representative motor and two of the sensory tests were chosen. There is no particular justification for the particular choice in each case, except for the fact that from the point of view of the experimental design they would tend themselves to fairly good agreement with the results (weakened) who were to be used. Other tests were also used (See Section 2.4) with smaller groups of patients, since excellent cooperation and achievement in all the tests was hardly to be expected.

1. Motor Tests

None of these tests were used in Cogan's study with patients.

However, there are some modifications here, in the manner of presentation

and in the methods of scoring² for which we found ample justification. The experimental period for the entire test was shortened in agreement with Cattell who considers the "....optimum length for an adult "p" test is one and one-half to two minutes...."³

The two motor tests were as follows:

a. Reverse-Stroke Z-test

After S is seated at the table, near E, he is asked the following question: "You know how to make a Z, like this (E writes letter Z similar to a printed one) don't you?" After the subject answers in the affirmative and sees E making the letter, he is given the pencil and is told, "Now, you make one like this."* After S has done as instructed, E continues the instruction as follows:

(1) "Now, when I say 'go,' start making as many of these letters as fast as you can, until I say 'stop.'" (Time allowed - 30 seconds). Then, E says: "That is fine, now, we shall make the letter Z the way it would appear in the mirror, like this (E writes letter down)." S follows instructions and writes letter down. Then, the following instructions are given:

(2) "Now, when I say 'go' start making these letters as fast as you can, until I say 'stop.'" (60 seconds). After this activity E proceeds by saying, "Now, we shall alternate these letters---first a regular Z, then a reversed one, like this." S writes down one part. The instructions then follow:

² See Section E, this Chapter.

³ H.B.Cattell, "On the Measurement of 'Perseveration,'" British Journal of Educational Psychology, 1935, 5, 90.

*See Figure 1, page 52.

and in the case of a single letter, the subject is justified.
The experimental period for the entire test was extended to 10 minutes.
The control and comparison tests were conducted for an equal test
one and one-half to two minutes.

The two major results were as follows:

1. Novice-Subject E-Test


After E is seated at the table, near E, he is asked the following question:
"You know how to make a E, like this E (writes letter E) similar to a
written one, don't you?" After the subject answers in the affirmative and
says E makes the letter, he is given the pencil and is told, "Now, you
make one like this." After E has done as instructed, E continues the
instruction as follows:

(1) "Now, when I say 'go,' start making as many of these letters
as fast as you can, until I say 'stop.'" (Time allowed - 30 seconds).
Then, E says: "That is fine, now, we shall make the letter E the way it
would appear in the mirror, like this E (writes letter down)." E follows
instructions and writes letter down. Then, the following instructions are
given:

(2) "Now, when I say 'go,' start making these letters as fast as you
can, until I say 'stop.'" (30 seconds). After this activity is completed by
saying, "Now, we shall alternate these letters--first a regular E, then
a reversed one, like this." E writes down one part. The instructions then
follow:

See Section E, this Chapter.
E. A. Gossard, "The Measurement of Psychomotor Skills,"
Journal of Educational Psychology, 1938, 3, 30.

FIGURE 1.
(Record of the Z-Test)



11. 20. 1911
(1911-12) 11. 20. 1911

11. 20. 1911

11. 20. 1911

(3) "When I say 'go,' start alternating them until I say 'stop.' Go as fast as you can." (30 seconds).

(4) "We shall now make regular Z's again, like the ones we did at first. Go as fast as you can. (30 seconds).

There are intervals of 10-15 seconds between each activity period. Thus, the four activity periods (3), 60, 30, 30 seconds) and the intervals last approximately 3-4 minutes altogether. Prolonged intervals are not desirable, since they may eliminate the effect of the perseverative tendency. Period (1) is only used as a "warming up" period, while the identical activity in (4) is actually used in the scoring.

b. The Grouped Stroke Test

The procedure in this test is quite similar to that in the "Z" test. There are four experimental periods as in the previous one. Each of these lasts 30 seconds. The intervals are the same as in the "Z" test. The instructions for the first activity are as follows:*

(1) "We shall now write three short strokes and a long one, like this, (E writes three short vertical strokes and a long vertical stroke that follows) then, "Now you make one like it. When I say 'go' you will start making many of them, one after the other, until I say 'stop.' (30 seconds).

The instructions for parts (2), (3), and (4) follow the same trend as those in the first motor test.

2. Sensory Tests

The first of these tests is, in principle, similar to the ones used by Wiersma, Jones and other investigators. We made use of some improvements

* See Figure 2, page 55.

(3) "When I say 'no', stop immediately even until I say 'stop'."

(4) "When I say 'no', stop immediately even until I say 'stop'."

(5) "When I say 'no', stop immediately even until I say 'stop'."

(6) "When I say 'no', stop immediately even until I say 'stop'."

(7) "When I say 'no', stop immediately even until I say 'stop'."

(8) "When I say 'no', stop immediately even until I say 'stop'."

(9) "When I say 'no', stop immediately even until I say 'stop'."

(10) "When I say 'no', stop immediately even until I say 'stop'."

(11) "When I say 'no', stop immediately even until I say 'stop'."

(12) "When I say 'no', stop immediately even until I say 'stop'."

2. General Instructions

The purpose of this test is to determine whether or not the subject can follow the instructions.

The subject is to be instructed to follow the instructions as given.

The subject is to be instructed to follow the instructions as given.

The instructions for the first activity are as follows:

(1) "When I say 'no', stop immediately even until I say 'stop'."

(2) "When I say 'no', stop immediately even until I say 'stop'."

(3) "When I say 'no', stop immediately even until I say 'stop'."

(4) "When I say 'no', stop immediately even until I say 'stop'."

(5) "When I say 'no', stop immediately even until I say 'stop'."

The instructions for parts (2), (3), and (4) follow the same pattern.

The instructions for the first activity are as follows:

3. General Instructions

The purpose of this test is to determine whether or not the subject can follow the instructions.

The subject is to be instructed to follow the instructions as given.

FIGURE 2.
(Record of the Stroke-Test)




PLATE 2.
(Record of the Gynae-Test)

22

in technique suggested by Shevach.⁴

The second test is, in some respects, an improvement on the methods previously used since the set-up of the machine (Polysine generator) is a very convenient means of gradating the stimuli.

a. Light Adaptation Test

A round hole (1.5 inches in diameter) was cut through a large, closed paper box, suspended from the ceiling. A 7.5 Watt, 115 volt General Electric bulb, wrapped in dark cloth was set up inside the box in a socket and connected with the house source of electric current. The hole which was high enough from the floor to be in the line of vision of an individual seated on a chair, was covered tightly with two sheets of typewriting paper. The subject was seated at a distance of four feet from the box. Suspended from the ceiling and above the line of vision of S was a 250 Watt bulb. The light was reflected upon the white paper covering the hole and was screened off on the other sides. The room was converted into a dark room.

After all the lights, but one, have been put out, the following instructions are given to S:

"Look straight ahead of you, on the white paper; I am going to put out the light. As soon as you see a round spot of light on the paper, you will tell me; the spot may not be very clear, but you will see it."

The lights are put out and E waits until S responds in the darkness. No measurements are taken at this point, since this adaptation is uncontrolled.

⁴ B.J. Shevach, "Studies in Perseveration VI. Methods for the Study of Sensory Perseveration," The Journal of Psychology, 1936, 3, 366.

is technique suggested by Sherrin.
The second test, in some respects, an improvement on the method previously used since it is set-up of the machine (the camera) is a very convenient means of obtaining the stimuli.

2. Light Adaptation Test

A round hole (1.5 inches in diameter) was cut through a large closed paper box, suspended from the ceiling. A 7.5 watt, 115 volt general electric bulb, wrapped in dark cloth was set up inside the box in a socket and connected with the house source of electric current. The hole which was high enough from the floor to be in the line of vision of an individual seated on a chair, was covered tightly with two sheets of transparent paper. The subject was seated at a distance of four feet from the box. Suspended from the ceiling and above the line of vision of the subject was a 250 watt bulb. The light was reflected upon the white paper covering the hole and was screened off on the other sides. The room was converted into a dark room.

After all the lights, etc. had been set out, the following instructions are given to the subject:
"Look straight ahead of you, on the white paper, I am going to turn out the light. I am going to have a round spot of light on the paper. You will tell me the spot was not so very clear, but you will see it. The lights are put out and I wait until your response in the darkness. No conversation is taken at this point, since this adaptation is necessary."

W. S. Hoar, "Changes in Sensation in Response to Light," The Journal of Psychology, 1906, 1, 250.

As soon as S responds, the 250 Watt light, only, is turned on for 60 seconds, the subject is, then, given the same instructions as above. The stop watch is started the moment the light is put out and is stopped the moment S responds by saying that he sees the "round spot of light." The lapsed time is the adaptation time. This procedure is repeated three times and three measurements are obtained.

b. Electrical Sensitivity Threshold

A MacIntosh⁵ Polysine Generator was used in determining the threshold for electrical stimulation of the skin. The subject is seated on a comfortable chair upon the arms of which the metal electrodes, to which the current from the machine is conducted, are attached. S places the index finger (beyond last joint) upon the electrodes on both sides of the chair. The dial of the machine is adjusted for an interrupted galvanic current (80 shocks per minute.) Before the switch is turned, S is instructed as follows:

"Put your fingers on here, like this (E helps S). Now, you will tell me as soon as you feel the slightest tingle in your fingers."

Then the current dial of the apparatus is slowly moved along the 0-180 milliampere scale until the instructed subject responds.

This procedure is repeated three or more times in order to get a consistent and reliable measure, which is the number of degrees traversed by the pointer of the dial on the milliampere scale.

3. A Test of Rhythmic Perseveration

In this test the process of hand-ear coordination is involved. A rubber tube closed on all sides and connected with a pneumatic recording

⁵ G.A. Remington, Gleanings of Low Voltage Techniques, 1930.

is shown as 2 seconds, the 200 watt light, only, is turned on for 20 seconds. The subject is, then, given the same instructions as above. The stop watch is started the moment the light is put out and is stopped the moment 2 responds by saying that he sees the "round spot of light." The elapsed time is the adaptation time. This procedure is repeated three times and three measurements are obtained.

2. Electrical Sensitivity Threshold

A Medinco[®] Polysine Generator was used in determining the threshold for electrical stimulation of the skin. The subject is seated on a comfortable chair upon the area of which the metal electrodes, to which the current from the machine is connected, are attached. 2 places the index finger (second last joint) upon the electrodes on both sides of the chair. The dial of the machine is adjusted for an interrupted galvanic current (50 shocks per minute). Before the switch is turned, 2 is instructed as follows:

"The four fingers on here, like this (2 holds 2), now, you will tell me as soon as you feel the slightest tingle in your fingers."

When the current dial of the apparatus is slowly moved along the 0-100 milliamperes scale until the instructed subject responds. This procedure is repeated three or more times in order to get a constant and reliable measure, which is the amount of current measured by the pointer of the dial on the milliamperes scale.

3. A Test of Spontaneous Polarization

In this test the process of back-ear coordination is involved. A rubber tube closed on all sides and connected with a pneumatic recording

set-up is presented to the subject. Only the piece of tubing is visible to the subject in front of the screen behind which the rest of the apparatus is to be found. The subject is instructed to press the tube once for every tick he hears. The metronome is then released and E shows the subject how to press it "as fast as the ticks are coming." S begins to press the tube, the stopwatch is started, and thus the actual experiment begins. The experimental period is divided into four parts.

(1.) Part one lasts for 30 seconds - from the subjects's first exerted pressure till 30 seconds have passed and, at that point the kymograph* is started.

(2.) Part two also lasts 30 seconds, after the kymograph runs for half a minute the electric key is pressed and the metronome speed is suddenly increased from 110 to 170 beats per minute.

(3.) Part three, with a fast metronome only, lasts for 15 seconds, since only the transfer from one speed to the other is the crucial point. At the end of the period, the metronome is stopped.

(4.) Period four is not really a part of the true experimental procedure, neither has it definite time limits. E observed that some patients keep on pressing the tube even after the metronome has stopped. Apparently some of the subjects become conditioned to the noise made by the kymograph which is not usually stopped simultaneously with the metronome. Some patients, however, continue pressing the tube even after both the metronome and the kymograph have been stopped. Moreover, a silenced kymograph does not eliminate these after-effects in some individuals.

During the first and second periods which last together for one

* See Kymographic records, p. 59.

set-up is presented to the subject. Only the phase of ending is visible to the subject in front of the screen behind which the rest of the apparatus is so designed. The subject is instructed to press the key once for every tick he hears. The metronome is then released and about the subject has to press it as fast as the ticks are coming. The key to press the tube, the stoppage is started, and thus the actual experimental period is divided into four parts.

- (1.) First one starts for 30 seconds - from the subject's time started pressure will 30 seconds after passed, and at that point the kymograph is started.
- (2.) Part two about lasts 30 seconds, after the kymograph runs for half a minute the electric key is pressed and the metronome speed is suddenly increased from 110 to 150 beats per minute.
- (3.) Part three, with a fast metronome only, lasts for 15 seconds, since only the transfer from one speed to the other is the crucial point at the end of the period, the metronome is stopped.
- (4.) Part four is originally a part of the first experimental procedure, neither one is definite time limit. It is provided that some patients have on pressing the tube even after the metronome has stopped. Apparently some of the subjects become so inclined to the end of the key the kymograph which is not usually stopped simultaneously with the metronome. Such patients, however, continue pressing the tube even after only the metronome and the kymograph have been stopped. Moreover, a finished kymograph does not eliminate these after-effects in some individuals.
- During the first and second periods which last together for one

FIGURE 3.

Kymographic Record of the Test of Rhythmic Perseveration

(A "No Change" Patient)



FIGURE 4.

Kymographic Record --- Rhythmic Perseveration

(A "Change" Patient)




FIGURE 3.

Hypnographic Record of the Test of Hypnotic Induction

(A "Change" Patient)

/

/

FIGURE 4.

Hypnographic Record --- Hypnotic Induction

(A "Change" Patient)

/

/

minute S gets used to a definite rhythmical activity (hand-ear) coordination at definite intervals. When the metronome speed is suddenly increased S "perseverates" and continues his previous activity for a small fraction of a minute until he completely adjusts himself to the new rhythm. The time utilized for the adjustment to the new rhythm and for the overcoming the tendency to preserve the old one is called the perseveration time. It is not measured in seconds, but in terms of the large waves traced from the moment of change on the kymograph which are shortly followed by the small waves which represent the new and faster rhythm.

D. Critique of Subjects' Reports (Reliability)

The report of psychotic subjects is not to be accepted naively, without any criticism. It must be stressed that E must make sure that S understands fully the instructions, but, this is not sufficient. It is necessary to focus S's attention upon the task (i.e., report et cetera) at hand. To some extent a tension and feeling of "need" to report must be created in the subject. In most cases it is fairly easy for E, after some experience, to identify the report, upon which the subject concentrated well. In the case of "adaptation to light" and "electrical sensitivity" a number of readings must be obtained, from which the reliability of the report can be deduced. When E felt that the value of the results of some test was impaired by lack of sufficient concentration on the part of S or by his excessive suggestibility, the results of that particular test were not used in the final computations. Thus, much of the work had to be discarded and consequently, the number of cases reduced, in accordance with this procedure. No unfair sampling was introduced as a result of this

minutes 2, used to a definite rhythmic activity (hand-part) corresponding to definite intervals. When the motoric speed is suddenly increased, B "perseverates" and continues his previous activity for a small fraction of a minute until he completely adjusts himself to the new rhythm. The time required for the adjustment to the new rhythm and for the overcoming the tendency to preserve the old one is called the perseveration time. It is not identical in subjects, but in terms of the large waves traced from the subject or change in the frequency which are shortly followed by the small waves which represent the new and faster rhythm.

D. Critique of Subjects' Reports (Reliability)

The report of psychomotor subjects is not to be accepted uncritically. Without any criticism, it must be stressed that B must make sure that B understands fully the instructions, but, this is not sufficient. It is necessary to insure B's attention upon the task (i.e., report of motor) at hand. To some extent a sensation and feeling of "need" to report must be created in the subject. In most cases it is fairly easy for B, after some experience, to identify the report, upon which the subject commented well. In the case of "distraction to light" and "reflexive sensitivity" a number of readings must be obtained, from which the reliability of the report can be deduced. When B fails to state the value of the results of some test was impaired by lack of sufficient concentration on the part of B or by his excessive suggestibility, the results of that particular test were not used in the final comparisons. Thus, much of the work had to be discarded and correspondingly, the number of cases reduced, in accordance with his procedure. No unit's analysis was introduced as a result of this

procedure, since there was no predominance of one diagnosis or another in the eliminated group. Suggestion played an important part in the experiment on electrical sensitivity when the subject felt a "tingle" even before the current was turned on. E. purposely asked for reports when there was no current passing in order to check the value of S's responses. A similar situation prevailed in the adaptation test, though it was more difficult to check the reliability of the reports.

In the other tests no such difficulties were noticed, since the results are dependent upon one's behavior and performance rather than "introspective" reports. No doubt, however, that even here, fluctuations in attention and in volition played an important and, at times, a vitiating role.

E. Methods of Scoring

The motor tests were scored, previously, in a number of different ways. The following are the different activities involved in each of the motor tests:*

X_1 - Habituated activity for 30 seconds (regular "Z's" (a) three short strokes and a long one).

Y - Different but similar activity for 60 seconds (reversed Z's and two short strokes and a long one).

Z - Alternation of habituated and different units for 30 seconds.

X_2 - Repetition of X_1 . Each activity in the first motor-test (2) has its parallel in the second motor test (grouped strokes).

Previous methods of scoring as reviewed by some investigators⁶

⁶ W. Stephenson, "An Introduction to So-Called Motor Perseveration Tests," British Journal of Educational Psychology, 1934, IV, 186-207.

*The letters X,Y,Z refer to the amount of output in each period.

procedures, since there was no presentation of one stimulus or another in the stimulus group. The stimulus group played an important part in the experiment of electrical sensitivity when the subject felt a "tingle" even before the current was turned on. It purposely varied the reports when there was no current passing in order to check the value of E's responses. A similar situation prevailed in the adaptation test, though it was more difficult to check the reliability of the reports.

In the other tests no such difficulties were involved, since the results are dependent upon one's behavior and performance rather than "introspective reports." No doubt, however, that even here, fluctuations in attention and in motivation played an important and, at times, a vital role.

Methods of Scoring

The motor tests were scored, previously, in a number of different ways. The following are the different activities involved in each of the motor tests:

- 1 - Habituated activity for 30 seconds (regular "E's") (a) three short strokes and a long one).
- 2 - Different but similar activity for 30 seconds (reversed E's and no short strokes and a long one).
- 3 - Alternation of habituated and different units for 30 seconds.
- 4 - Repetition of A. Each activity in the first motor-test (1) was repeated in the second motor test (grouped strokes).

Previous methods of scoring as reviewed by some investigators

W. F. Floyd, "An Introduction to the Study of Human Behavior," 1928, IV, 146-157.

*The letters A, Y, Z refer to the amount of activity in each period.

possessed many masking factors. For instance, a scoring equivalent to $X + Y - 2Z$ will give too much weight, in the "p" score, to the speed in the performance of the subject. Also, $\frac{2X}{YZ}$ gives undue weight to the "different but similar activity" in which the difficulty is more of learning the new mode of response, rather than in the interference of perseveration.

The rationale of the present scoring follows a different path. The X_1 activity is considered as a "warming up" activity, not be counted in calculating the final score. Then, in Z-activity (alternation), if there were no perseveration at all, the number of letters written would be equal to half the number written in X_2 plus one fourth the number written in Y (since there a 60 second period is allowed). In other words, there being no perseveration $Z = \frac{X}{2} + \frac{Y}{4}$ or $2Z = X + \frac{Y}{2}$.

However, Z is usually smaller on account of "p" in the alternating activity; then, the number of units lost due to perseveration will equal to

$$X + \frac{Y}{2} - 2Z \text{ or } 2X + Y - 4Z.$$

However, this alone will give us a quantity, not related to the speed of the individual will be the adequate "p" score. Hence we have

$$\frac{X + \frac{Y}{2} - 2Z}{X + \frac{Y}{2}} (100) \text{ as the "p" score; i.e., the ratio between the units}$$

(letters or groups of strokes) lost due to "interference." This method of scoring was employed in the preliminary results present in the present study. It is noteworthy that when this method is used, negative scores⁷ (due to a larger 2Z value than $X + \frac{Y}{2}$) are obtained. Such scoring is undesirable for a number of statistical and theoretical reasons.

A different method of scoring was applied to the results of the

⁷ See Chapter VI, present study.

assesses many masking factors. For instance, a scoring equivalent to $X + Y - 2Z$ will give too much weight, in the "p" score, to the speed in the performance of the subject. Also, $\frac{2X}{Y}$ gives undue weight to the "difference of similar activity" in which the difficulty is more of learning the new code of response, rather than in the interference of perseveration.

The rationale of the present study follows a different path. The

activity is considered as a "learning up" activity, not to be counted in evaluating final score. Then, in 2-activity (perseveration), if there are no perseveration at all, the number of letters written would be equal to half the number written in X plus one fourth the number written in Y since there is 80 second period is allowed. In other words, these being

$$\text{perseveration } Z = \frac{Y}{2} + \frac{X}{4} \text{ or } 2Z = X + \frac{Y}{2}$$

however, Z is usually smaller on account of "p" in the perseverating activity

then, the number of letters lost due to perseveration will equal to

$$\frac{X}{2} - Z \text{ or } 2X + Y - 4Z$$

however, this alone will give us a quantity, not related to the speed of

the individual will be the separate "p" score. Hence we have

$$+ \frac{Y}{2} - \frac{2Z}{X} (100) \text{ as the "p" score; i.e., the ratio between the units}$$

letters or groups of strokes) lost due to "interference." This method of

curious was employed in the preliminary results present in the present study

is noteworthy that when this method is used, negative scores (due to a

greater 2Z value than $X + \frac{Y}{2}$) are obtained. Such scoring is undesirable for

number of statistical and theoretical reasons.

A different method of scoring was applied to the results of the

final study. In order to deal largely with positive scores and provide for simpler calculations, an output ratio is set up in which the habitual regular output is put over the alternating activity which is to reveal perseveration. The ratio is as follows: $\frac{X+Y}{2Z}$. If the person is a great perseverator he will have many fewer units in his "Z" output than in either "X" or "Y"; thus, the ratio will be larger; consequently, the "p" score will also be larger. This ratio is multiplied by 100 in order to avoid dealing with fractions in the frequency distribution and in the conclusions.

Scoring the adaptation test was quite simple---the mean of the last two measures of time necessary to see the "round spot" was used as the final score. The assumption here is, as it has been in previous experiments,⁸ that the longer the after-function or after-sensitivity, the greater is the "Sekundaerfunction" (secondary function) or perseveration.

The strength of current necessary to "just feel a tingle" was taken also as a measure of the subject's perseveration. Here too, the average of the last two readings was taken as the final score. The basic idea behind this is that inertia is a property of states of rest as well as of motion; the higher perseverator would have a higher threshold for any sensitivity. His inertia (i.e., perseveration) is that of maintaining a state of rest despite disturbances. However, when stimulation is strong enough, this "inertia" is overcome, the subject then "feels" the stimulation. The point at which the liminal stimulation begins is a measure of perseveration, since up to that point the inertia or perseveration of rest is maintained.

In the case of "rhythmic perseveration," the time that lapses until

⁸ See Chapter II, present study.

final study. In order to deal largely with positive scores and provide for simpler calculations, an output ratio is set up in which the habitual output is put over the alternating activity which is to reveal perseveration. The ratio is as follows: $\frac{X+Y}{2}$. If the person is a great perseverator he will have many fewer units in his "X" output than in either "Y" or "Y'"; thus, the ratio will be large; consequently, the "p" score will also be large. This ratio is multiplied by 100 in order to avoid dealing with fractions in the frequency distribution and in the conclusions. Scoring the adaptation test was quite simple---the mean of the last two measures of time necessary to see the "round spot" was used as the final score. The assumption here is, as it has been in previous experiments, that the longer the after-function or after-sensitivity, the greater is the "secondary function" (secondary function) or perseveration. The strength of current necessary to "just feel a tingle" was taken also as a measure of the subject's perseveration. Here too, the average of the last two readings was taken as the final score. The basic idea behind this is that inertia is a property of states of rest as well as of motion; the higher perseverator would have a higher threshold for any activity. His inertia (i.e., perseveration) is that of maintaining a state of rest despite disturbances. However, when stimulation is strong enough, this "inertia" is overcome, the subject then "feels" the stimulation. The point at which the minimal stimulation begins is a measure of perseveration, since up to that point the inertia or perseveration of rest is maintained. In the case of "rhythmic perseveration," the time that lapses until

adjustment to the new rhythm of the metronome takes place is considered as a measure of perseveration. However, that time cannot be measured in any other manner except in terms of distance traversed on the smoked drum of the kymograph. Every pressure exerted by S on the tube, records a long wave on the kymograph drum. The waves of the first 30 seconds are somewhat larger (slower metronome) than the ones in the following 15 seconds (faster metronome). The smaller waves appear shortly after the speed has been changed. A few large waves still persist. Hence E measures by means of dividers the length of the first five waves (apt to include the few persisting from the previous rhythm) and then, the length of the last five waves in the 15 second period. The ratio between the former (supposed to be larger) and the latter gives an index of perseveration.

The final score is calculated in the following manner:

$$\frac{L_1}{L_2} \times 100 = P.$$

L_1 is the length of the first five waves immediately after the change of speed of the metronome; L_2 is the length of the last five waves of that fast metronome 15 second period---P is the final score.

adjustment to the new rhythm of the metronome takes place is considered as a measure of perseveration. However, that time cannot be measured in any other manner except in terms of distance traversed on the smoked drum of the kymograph. Every pressure exerted by S on the tube, records a line on the kymograph drum. The waves of the first 30 seconds are somewhat larger (lower metronome) than the ones in the following 30 seconds (faster metronome). The smaller waves appear shortly after the speed has been changed. A few large waves still persist. Hence E measures by means of the length of the first five waves (not to include the few waves from the previous rhythm) and then, the length of the last five waves in the second period. The ratio between the former, supposed to be larger, and the latter gives an index of perseveration.

The final score is calculated in the following manner:

$$\frac{I_1}{I_2} \times 100 = P.$$

I_1 is the length of the first five waves immediately after the change of speed of the metronome; I_2 is the length of the last five waves of that second metronome in second period--P is the final score.

4. Preliminary Results

The aim of this study was much more circumscribed during the preliminary period of experimentation. I was interested in determining the difference between the two groups and the intercorrelation of the tests employed. The picture presented itself in a somewhat different than it appeared in other investigations. The table below gives us the means of the scores of four tests* for the two diagnostic groups.

CHAPTER V

TABLE VI
EXPERIMENTAL RESULTS

Mean T-Scores in Cases of Depressive Reaction and Manic-Depressive

Diagnosis	T-test	Percept	Alph. Digit.	Alph. Digit.	Age
Depressive Reaction	23.1	-2.0	22.9	12.5	23.4
Manic-Depressive	24.3	13.1	22.0	20.3	23.1
S_{xy}/S_{xx}	.37	31.2	3.55	1.74	
Chance is 100	27	25	25	25	

There appears to be a definite trend, on the basis of these data, towards higher preservation among the depressives and lower among the manic-depressives. Of course, in case of the T-test and the association test, the differences appear very small and insignificant. However, in the other tests they are of considerable magnitude. Another interesting fact was simultaneously revealed: the mean age of the

*Also the "fig" test was used; for little data presented the language

CHAPTER V

EXPERIMENTAL RESULTS

A. Preliminary Results

The aim of the study was much more circumscribed during the preliminary period of experimentation. I was interested in determining the difference between the two groups and the intercorrelation of the tests employed. The picture presented itself in a manner somewhat different than it appeared in other investigations. The table below gives us the means of the scores of four tests* for the two diagnostic groups.

TABLE VI

mean P-Scores in Cases of Dementia Praecox and Manic-Depressive

Diagnosis	Z-test	Strokes	Elect. Sensit.	Adapt.	Age
Dementia Praecox	23.1	-2.0	56.9	15.0	42.4
Manic Depressive	24.2	15.1	63.0	20.2	33.1
D/FE_D	.27	31.9	2.50	1.24	
Chances in 100	57	98	95	80	

There appears to be a definite trend, on the basis of these data, towards higher perseveration among the depressives and lower among the praecoxes. Of course, in case of the Z-test and the adaptation test, the differences appear very small and insignificant. However, in the other tests they are of considerable magnitude (grouped strokes, stimulation). Another interesting fact was simultaneously revealed; the mean age of the

*Also the "it" test was used; too little data prevented its inclusion.

A. Preliminary Results

The aim of the study was much more circumscribed during the preliminary period of experimentation. I was interested in determining the difference between the two groups and the interpretation of the tests employed. The picture presented itself in a manner somewhat different than it appeared in other investigations. The table below gives the means of the scores of four tests* for the two diagnostic groups.

TABLE VI

Mean T-Scores in Cases of Dementia Precox and Manic-Depressive

Diagnosis	T-Test	Stroop	Block Design	Adapt. Test	Age
Dementia Precox	23.1	25.0	20.9	15.0	42.4
Manic-Depressive	24.8	22.1	22.0	20.2	32.1
Age	27	21.2	22.0	1.24	
Stroop in	27	28	22	20	

There appears to be a definite trend, on the basis of these data, towards higher perseveration among the depressives and lower among the manics. Of course, in case of the T-test and the adaptation test, the differences appear very small and insignificant. However, in the other tests they are of considerable magnitude (grouped strokes, stimulation). Another interesting fact was simultaneously revealed; the mean age of the "Also the 'it' test was used; too little data prevented its inclusion.

of the praecox group was nine years higher than that of the manic-depressive group. The question arose, therefore, whether the differences between the means of the two groups are actually due to the differential diagnosis or to a possible correlation, negative or positive, between the tests and the age of the subjects. All the tests were, therefore, correlated with age. The rank-difference method was used in all correlations of the preliminary results.

The table below shows us the correlations of each test with age. Among these coefficients of correlation, the first one (adaptation) is the only significant one; its probable error being .099, less than one fourth its own magnitude.

TABLE VII

Preliminary Correlations of Tests with Age.

Light Adaptation	Electrical Stimulation	Z-Test	Strokes
.43	.27	-.26	-.31

The other coefficients of correlation are much less than four times their probable errors. Hence, the only significant correlation is a positive one, and if anything, it would reduce still more the adaptation score in the case of the dementia praecox group, taking age into account, and thus, accordingly increase the difference between the latter group and that of manic-depressive.

The intercorrelation of the tests presented a case for the separation of the motor from the sensory tests. The correlation between the two motor

The present group was nine years higher than that of the manic-depressive group. The question arises, therefore, whether the difference between the two groups is actually due to the differential diagnosis or to a possible correlation, negative or positive, between the tests and the age of the subjects. All the tests were, therefore, correlated with age. The rank-difference method was used in all correlations of the preliminary tests.

The table below shows the correlations of each test with age. Only three coefficients of correlation, the first one (adaptation) is only significant one; the probability error being .032, less than one with the two negative.

TABLE VII

Preliminary Correlations of Tests with Age.

Adaptation	Electrical Stimulation	E-Test	Stroop
.43	.37	-.36	-.21

Other coefficients of correlation are much less than four times their probable error. Hence, the only significant correlation is a positive one. If anything, it would reduce still more the adaptation score in the case of the manic-depressive group taking age into account, and that, accordingly, across the difference between the latter group and that of manic-depressive. The interpretation of the tests presented a case for the adaptation the better from the present tests. The correlation between the two major

tests (Z and grouped stroke) was a rather significant one; it was a positive correlation of .51 with a P. E. of .095. Thus it appears that there definitely is a linkage between the two tests measuring a like factor.

The correlation between the adaptation test and that of electrical liminal sensitivity was positive, but rather low (.27) and not quite four times its P. E. One may thus presume a weak factor of sensory perseveration.

A more important discovery, however, was the negligible relationship between the motor and sensory tests. For instance, the positive coefficient of correlation between the Z-test and that of adaptation was .01. It was, therefore, necessary to conclude, tentatively, on the basis of these results, that the sensory tests used may not measure perseveration at all. They may measure the reaction-time of the subjects or their ability to follow instructions, rather than the perseverative tendency. Or, if there are such things as motor and sensory perseveration, some relationship between them ought to be shown in order to permit the common term.

The following problems suggested themselves for further investigation with a larger number of subjects:

1. Is there a difference in perseveration between like age groups of different psychiatric diagnosis?
2. Do we still discern two distinct factors, i.e., motor and sensory perseveration?
3. What age level is responsible for the correlation with age (if present)?

The results in the next section are attempts to solve these problems.

tests (S and grouped strokes) was a rather significant one; it was a positive correlation of .51 with a P. E. of .003. Thus it appears that there finally is a linkage between the two tests measuring a like factor. The correlation between the adaptation test and that of electrical manual sensitivity was positive, but rather low (.23) and not quite four times the P. E. One may thus presume a weak factor of sensory perseveration. A more important discovery, however, was the negligible relationship between the motor and sensory tests. For instance, the positive coefficient of correlation between the E-test and that of adaptation was .01. It was, therefore, necessary to conclude, tentatively, on the basis of these results, that the sensory tests used may not measure perseveration at all. They may measure the reaction-time of the subjects or their ability to follow instructions, rather than the perseverative tendency. Or, if there is such things as motor and sensory perseveration, some relationship between them ought to be shown in order to permit the common term.

The following problems suggested themselves for further investigation with a larger number of subjects:

1. Is there a difference in perseveration between like age groups of different psychiatric diagnoses?
2. Do we still discern two distinct factors, i.e., motor and sensory perseveration?
3. What age level is responsible for the correlation with age (if present)?

The results in the next section are attempts to solve these problems.

B. Further Results

The tables (VIII, IX, X, XI, on pages 69, 70, 71, 72) are compiled on the basis of the results obtained from the second experimental group (B).

Table VIII shows, as far as age is concerned, the reverse condition from what is obtained in Table VI. Here the mean age of the manic-depressives is higher than that of the dementia praecox cases. The other conditions, however, remain approximately the same, that is, the scores of the dementia praecox patients in all tests are lower than those of the manic-depressives. A truly significant difference exists, perhaps, in the case of the Z-test (91 chances in 100); the reliability of the other differences, though pointing in one direction, is not great enough to serve as a basis for certainty and conclusiveness. Of course, in both cases (of preliminary and present results) only the depressive-phase patients were used in the comparison.

Tables IX, X, and XI give us the further opportunity to analyze the scores within the general praecox group; groups of patients of the same number, whose mean age was approximately the same, were matched. There, the catatonic group, seems to be definitely higher in three of the tests, than either group of paranoid praecoxes and the "others" (hebephrenics and simple) group. This was further substantiated (though not perfectly) after the significant ratios had been computed. The differences between the means of the catatonic group and each of the others is of considerable significance, at least in two tests (Strokes and adaptation). Obviously, the differences between the paranoid and "others" group is of no significance. The dementia praecox paranoid group seems to be lowest in the general praecox group in regard to the adaptation and electrical sensitivity tests. Otherwise, there is little difference between the two groups (paranoid and others).

The tables (VIII, IX, X, XI, on pages 52, 53, 54, 55) are arranged on the basis of the results obtained from the second experimental group (5). Table VII shows, as far as age is concerned, the reverse condition. The data is contained in Table VI. Data for mean age of the manic-depressive is higher than that of the dementia praecox cases. The other condition, however, results approximately the same, that is, the scores of the dementia praecox patients in all tests are lower than those of the manic-depressive. A fairly significant difference exists, perhaps, in the test of the 2-test (41 chances in 100); the reliability of the other differences, though pointing in one direction, is not great enough to serve as a basis for certainty and conclusiveness. Of course, in both cases (of manic-depressive and dementia praecox) only the depressive-phase patients were used in the comparison.

Tables IX, X, and XI give us the further opportunity to analyze the scores within the general praecox group; groups of patients of the same age, whose mean age was approximately the same, were matched. Thus, the manic group, seems to be definitely higher in three of the tests, than the group of dementia praecox and the "other" (schizophrenic and manic) group. This was further substantiated (though not perfectly) at the significant test had been computed. The difference between the mean of the manic group and each of the others is of considerable significance, at least in two tests (Schizophrenic and manic). Obviously, the difference between the manic and "other" group is of no significance. The dementia praecox group seems to be lower in the general praecox group in regard to the agitation and electrical sensitivity tests. Otherwise, there is little difference between the two groups (manic and other).

TABLE VIII

MEANS OF P SCORES FOR DEMENTIA PRAECOX AND MANIC-DEPRESSIVE CASES

	Dementia Praecox	Manic Depressives	$\frac{D}{PE_D}$	CHANCES IN 100*
Age	35.2	40.2		
Z-Test	116.5	130.5	2.00	91
Strokes	110.8	114.7	.86	72
Adaptation	44.9	51.4	1.16	78
Electrical Stimulation	61.8	64.7	1.02	75

*Number of chances in 100 that the obtained difference is significant (that a true difference exists).

TABLE VIII

MEANS OF P-VALUES FOR ORBITAL PHASES AND MANIC-DEPRESSIVE STATES

Age	Depressive Phase	Manic Depressive	$\frac{D}{D+M}$	CHANGES IN 100*
20-25	50.2	40.2		
26-30	116.5	180.5	5.00	31
31-35	112.8	114.7	.68	75
36-40	44.9	51.4	1.15	76
41-45	61.8	64.7	1.05	75

*Number of changes in 100 that the obtained difference is significant
(based on difference table).

TABLE IX

P-Scores of Catatonics and other Praecox Cases

	Catatonics	Others	$\frac{D}{PE_D}$	Chances in 100
Age	35.7	36.1		
Z-Test	124.7	112.3	1.69	90
Stroke	129.7	111	2.93	97
Adaptation	60.6	36.6	3.58	99
El. St.	66.	60.3	1.60	86

TABLE IX

I-Scores of Catalonics and Other Protein Cases

	Catalonics	Others	$\frac{D}{P_{50}}$	Changes in 100
Age	33.7	35.1		
2-Test	124.7	117.3	1.53	30
Stroke	123.7	111	2.33	27
Abstinence	60.3	38.3	3.58	33
El. St.	58	60.3	1.50	33

TABLE X

P-Scores of Catatonics and Paranoiacs

	Catatonics	Paranoiacs	$\frac{D}{PE_D}$	Chances in 100
Age	35.7	35.7		
Z-Test	124.7	117.7	1.10	77
Stroke	129.7	110.7	3.16	98
Adaptation	60.6	31.1	4.35	100
El.St.	68.0	60.3	1.52	84

TABLE I

Properties of Catalysts and Parameters

Catalyst	Catalytic Activity		Temperature, °C	Pressure, mm Hg
	Rate, %/hr	Yield, %		
1	10.0	10.0	100	100
2	12.5	12.5	110	110
3	15.0	15.0	120	120
4	17.5	17.5	130	130
5	20.0	20.0	140	140

TABLE XI

P-Scores of Paranoiacs and "Others"

	Paranoiacs	Others	$\frac{D}{PE_D}$	Chances in 100
Age	35.7	36.1		
Z-Test	117.7	112.3	1.33	62
Strokes	110.7	111	.08	52
Adaptation	31.1	36.6	1.55	65
El. St.	60.3	60.3		50

TABLE XI

T-Geometries of Enantiomers and "Others"

Enantiomers in 100	$\frac{D}{P}$	Others	Enantiomers	
		50.1	50.7	Ag
43	1.50	50.5	51.7	2-Test
50	.75	51.1	51.7	Strocas
53	1.50	50.5	51.1	Adaptation
50		50.5	50.5	El. St.

In Table XII on page 74, the manics are compared with the depressives, there are no significant differences in the scores of the Stroke-test and that of electrical sensitivity, but the differences on the other two tests are of considerable magnitude and the chances are 99 in 100 that a true difference exists.

The next table (XIII), on page 75, gives a complete picture of the intercorrelations* of the various tests with each other and with age. Except for the test of electrical sensitivity (in this case the correlation is significant), age plays a small role in modifying the "p" score of individuals. There is a slight correlation with age between each of the tests. The next two highest positive correlations are between the two sensory and between the two motor tests respectively. One of the motor tests correlates very slightly, positively with each of the sensory ones, but the other motor tests (strokes) shows a negative coefficient of correlation in both cases.

The results in Tables XIV, XV, on page 76 are rather unexpected. It appears that the scores of the three better intercorrelated tests are much lower for the lowest age group (20-29). The scores then go up and stay up between 30-49 and then are reduced again, at the highest age level (50-59). The differences between the 20-29 and 30-39 groups are quite reliable and significant, showing a definite trend upward, beyond the 30 year level. The reduction of the scores at the highest age-level (50-59) is less marked and not quite so significant.

The table below, (XVI) gives the mean results obtained for a group of partly recovered patients. The improvement was determined by the subject's

*The Product-moment method of correlation was used.

In Table XII on page 74, the means are compared with the
 themselves, there are no significant differences in the scores of the
 one-test and that of electrical sensitivity, but the differences on the
 other two tests are of considerable magnitude and the chances are 99 in 100
 that a true difference exists.

The next table (XIII), on page 75, gives a complete picture of the
 "correlations" of the various tests with each other and with age.
 Except for the test of electrical sensitivity (in this case the correlation
 is significant), age plays a small role in modifying the "u" scores of
 individuals. There is a slight correlation with age between each of the
 tests. The next two highest positive correlations are between the two
 sensory and between the two motor tests respectively. One of the motor
 tests correlates very slightly, positively with each of the sensory ones,
 and the other motor tests (stroked) shows a negative coefficient of
 correlation in both cases.

The results in Tables XIV, XV, on page 76 are rather unexpected. It
 seems that the scores of the three better intercorrelated tests are much
 lower for the lowest age group (20-29). The scores then go up and stay up
 between 30-39 and then are reduced again, at the highest age level (50-59).
 The differences between the 20-29 and 30-39 groups are quite reliable and
 significant, showing a definite trend upward, beyond the 39 year level.
 A reduction of the scores at the highest age-level (50-59) is less marked
 and not quite so significant.

The table below (XVI) gives the mean results obtained for a group of
 fully recovered patients. The improvement was determined by the subject's

TABLE XII

A COMPARISON BETWEEN MANIA AND DEPRESSION IN REGARD TO PERSEVERATION

	Manics	Cases of Depression	$\frac{D}{PE_D}$	Chances in 100
Age	34.5	35.5		
Z-Test	119.3	152.3	3.55	99
Stroke	109	107.8	.29	58
Adaptation	39.3	64.9	3.46	99
Electrical Stimulation	65.1	58.8	1.26	60

TABLE III

A COMPARISON BETWEEN NINA AND DEEPWATER IN REGARD TO PERCENTAGE

	Percent of Depression	$\frac{N}{P}$	Changes in 100
Age	84.8		
W-Tot	81.3	3.55	35
Stress	108	.73	36
Adaptation	83.3	3.16	39
Electrical stimulation	68.1	1.76	60

TABLE XIII

CORRELATIONS BETWEEN THE TESTS

	Age	Adapt. to Light	Elect. Sensit.	Z-test	Stroke Test
Age		.122 $\pm .062$	.348 $\pm .079$	.284 $\pm .063$	.113
Adaptation to Light	.122 $\pm .062$		.438 $\pm .071$	.185 $\pm .069$	-.201
Electrical Sensitivity	.358 $\pm .079$	.438 -.071		.096 -.090	-.209
Z-test	.284 $\pm .063$	.185 -.069	.096 $\pm .090$		.306 $\pm .065$
Stroke-test	.113	-.201	-.209	.306 $\pm .065$	

TABLE XIV

TEST SCORES FOR YOUNGER AGE-GROUPS

	20-29	30-39	$\frac{D}{PE_D}$	Chances in 100
Z-Test	115.4	127.9	3.90	100
Grouped Strokes	115.4	102.2	2.94	97
Adaptation	34.0	48.2	3.71	99
El. Stim.	52.9	63.2	2.69	97

TABLE XV

TEST SCORES FOR OLDER AGE-GROUPS

	40-49	50-59	$\frac{D}{PE_D}$	Chances in 100
Z-Test	131.1	118	2.27	94
Grouped Strokes	123.0	110	2.57	95
Adaptation	49.4	39.8	2.73	97
El. Stim.	64.0	63.7	.09	52

TABLE IV

TEST RESULTS FOR VARIOUS AGE GROUPS

Age Group	40-49	50-59	$\frac{D}{FSD}$	Changes in 100
4-7 test	110.4	107.5	2.99	100
Propped strokes	110.4	107.5	2.99	97
Substitution	107.0	107.5	2.91	97
El. Sign.	107.0	107.5	2.99	97

TABLE IV

TEST RESULTS FOR OTHER AGE GROUPS

Age Group	40-49	50-59	$\frac{D}{FSD}$	Changes in 100
4-7 test	110.1	110	2.92	94
Propped strokes	110.0	110	2.97	92
Substitution	107.0	107.0	2.92	97
El. Sign.	107.0	107.0	2.92	92

TABLE XVI

P-Scores of Improved Subjects

Z-Test	Stroke	Adaptation	Electrical Stimulation
120	118	37.7	63.9

accessibility as determined by E in cooperation with the attending physician. Some of this group were on parole which is the status nearest to dismissal from the institution.

The only test that shows any definite difference is that of adaptation time. The improved patients present in this one test a much smaller score. In the rest of the tests no great change in "p" is evident when compared with the mean of the entire experimental group.

C. The Test on Rhythmic Perseveration

The subjects were divided into two groups after the results were obtained. Some of the subjects demonstrated no change in their speed of movement (pressing the tube) after the speed of the metronome was altered. The other group showed definite changes in the rhythm shortly after the metronome beats came more rapidly. The individual differences within this latter group are not sufficiently marked nor can they be adequately graded in order to be correlated with the other tests. However, there are some differences between the magnitude of the scores of the subjects who demonstrate no change in rhythm and those who do show the change. In Table XVII (below) we see that in three of the other tests (except adaptation)

TABLE XVI

F-Scores of Improved Subjects

2-Test	Errors	Adaptation	Electrical Stimulation
120	118	27.7	22.2

cessibility as determined by 2 in cooperation with the attending physician. Some of this group were on parole which is the status nearest to dismissal from the institution.

The only test that shows any definite difference is that of adaptation time. The improved patients present in this one test a much better score. In the rest of the tests no great change in "p" is evident when compared with the mean of the entire experimental group.

The Test on Rhythmic Perseveration

The subjects were divided into two groups after the results were obtained. Some of the subjects demonstrated no change in their speed of movement (pressing the tube) after the speed of the metronome was altered. Another group showed definite changes in the rhythm shortly after the metronome beats came more rapidly. The individual differences within this latter group are not sufficiently marked nor can they be adequately graded in order to be correlated with the other tests. However, there are some

differences between the magnitude of the scores of the subjects who demonstrate no change in rhythm and those who do show the change. In Table II (below) we see that in three of the other tests (except adaptation)

TABLE XVII

P-Scores for High and Low Perseverators as
Determined by the Test on Rhythmic Perseveration.

No Change				Change			
Z-Test	Strokes	Adapt.	Elect. Stimul.	Z-Test	Strokes	Adapt.	Elect. Stimul.
128	114	41.6	65.3	120	110	45.3	59.7

the mean scores for the "no change" group are higher than for the "change" group.

Much more should be expected from this method with an electrical set-up of more exact recording devices.

TABLE XVII

P-Scores for High and Low Frequency Groups as
Determined by the Test on Rhythmic Perception.

No Change				Change			
2-Test	Strokes	Adapt.	2-Test	Strokes	Adapt.	2-Test	Strokes
125	114	41.8	55.2	125	110	45.8	59.7

The mean scores for the "no change" group are higher than for the "change"

group.

Which more should be expected from this method with an electrical

set-up of more exact recording devices.

2. Depressive and S-Depressive

Our results seem to agree in part with the tables presented by Stephenson.² In his results Stephenson established the fact that mania are lowest performers, while next to them, somewhat higher, are some cases of depressive psychosis (see Table I). Then with higher percentages are the manic-depressives, while the highest performers are, according to him, a number of depressive cases (Table II). Our results appear to confirm the fact that the mania are lowest performers (with the exception of the test on electrical excitability). Next it appears that the manic-depressive group as a whole is higher than the depressive group in every test, both in the preliminary as well as in the final readings. The mean age of the manic-depressive's being higher in the preliminary results and lower in the final ones. This is also quite a difference from that obtained by Shipley³ who gives the manic-depressive group highest in regard to perseveration. The reason for the obvious inconsistency may be found in the following explanation:

DISCUSSION OF RESULTS

1. The methods were quite different in Shipley's and in Stephenson's work. The former used entirely different tests (psychomotor, verbal, excitability, attention, and memory). The latter's manner of obtaining the scores on the tests of a 50 point scale may have changed the true picture to some extent. No measure of the significance of the difference between the mean was employed by either investigator.

2. No differentiations have been made within the manic group. There may be types of individuals in this group who may behave the same up to the rest of the group.

¹ See Table III, page 56.

² See Table IV, page 56.

DISCUSSION OF RESULTS

CHAPTER VI

A. Diagnoses and P-Scores

Our results seem to agree in part with the tables presented by Stephenson.¹ In his results Stephenson establishes the fact that manics are lowest perseverators, while next to them, somewhat higher, are some cases of dementia praecox (low "p"), then still higher perseverators are the manic-depressives, while the highest perseverators are, according to him, a number of praecox cases (high "p"). Our results appear to confirm the fact that the manics are lowest perseverators (with the exception of the test on electrical sensitivity). Next it appears that the manic-depressive group as a whole is consistently higher in every test, both in the preliminary as well as in the final results; the mean age of the manic-depressive's being higher in the preliminary results and lower in the final ones. This is also quite a different picture from that obtained by Shipley² who places the schizophrenic group highest in regard to perseveration. The answer to the obvious inconsistency must be found in the following explanations:

1. The methods were quite different in Shipley's and in Stephenson's work. The former used entirely different tests (psychogalvanic reflex, conditionability, extinction, and others). The latter's manner of combining the scores on the basis of a 20 point scale may have changed the true picture to some extent. No measures of the significance of the difference between the means were computed by either investigator.

2. No differentiations have been made within the dementia praecox group. There may be types of individuals in this group who may bring the score up for the rest of the group.

¹ See Table III, page 37.

² See Table IV, page 40.

Our results seem to agree in part with the tables presented by Johnson.¹ In his results Johnson established the fact that manic-depressives, while next to them, somewhat higher, are some cases of dementia praecox (low "q"), than still higher parasympathetic are manic-depressives, while the highest parasympathetic are, according to him, a number of parasympathetic cases (high "q"). Our results appear to confirm the fact that the manic-depressives (with the exception of the low of electrical negativity). Next it appears that the manic-depressive group as a whole is consistently higher in every test, both in the preliminary as well as in the final results. The mean age of the manic-depressive's being higher in the preliminary results and lower in the final ones. This is also quite a different picture from that obtained by Shiley,² who gives the schizophrenic group highest in regard to parasympathetic. The answer to the obvious inconsistency must be found in the following explanations:

1. The methods were quite different in Shiley's and in Johnson's work. The former used entirely different tests (psychodynamic, etc., sensibility, extinction, and others). The latter's manner of obtaining the scores on the basis of a 50 point scale may have changed the true picture to some extent. No measures of the significance of the differences between the means were computed by either investigator.
2. No differentiations have been made within the dementia praecox group. There may be types of individuals in this group who may differ from one another for the rest of the group.

¹ See Table III, page 37.
² See Table IV, page 30.

When we sought to try and distinguish the praecox cases on the basis of the degree of dementia³ we found that too subjective criteria will have to be used. Hence the more "objective" groupings used in psychiatry were employed. The results show (again, with the exception of electrical stimulation) that within the praecox group, the catatonics show a markedly higher degree of perseveration than do the other two praecox groups. We tried to eliminate the age factor in this comparison in order to make it much more correct. These results, then, may explain Stephenson's "high-p" cases and "low-p" cases. The former may correspond to our catatonic group, while the latter may be characteristic of the paranoid, hebephrenic and simple types of dementia praecox.

Parenthetically, it should be remarked that even the catatonic group does not exceed in all tests the mean score of the depressive-phase psychotics. The results concerning the catatonics seem to be in agreement with those obtained by Mays⁴ in his studies.

Therefore, if we were to average up the scores we would be able to show the following sequence in order of increasing perseveration.

1. Manic
2. Dementia praecox paranoids
3. Hebephrenic and simple praecox patients
4. Manic-depressives
5. Catatonics

³ W. Stephenson, "Studies in Experimental Psychiatry 11. Some Contacts of p.-factor in Psychiatry," Journal of Mental Science, 1932, 76, 315.

⁴ L.L. Mays, "Studies in Catatonia V. Perseverational Tendencies in Catatonic Patients," The Psychiatric Quarterly, 1934, 8, 728.

When we sought to try and distinguish the present cases on the basis of the degree of dominance, we found that too subjective criteria have to be used. Hence the more "objective" groupings used in the study were employed. The results show (again, with the exception of the clinical situation) that within the present group, the catatonic show markedly higher degrees of perseveration than do the other two present groups. We tried to eliminate the age factor in this comparison in order to make it much more correct. These results, then, may explain Stephenson's "high" cases and "low" cases. The former are comparable to our catatonic group, while the latter may be comparable to the paranoid, schizophrenic and simple types of dementia praecox. Paradoxically, it should be remembered that even the catatonic group does not exceed in all tests the mean score of the depressive group. The results concerning the catatonics seem to be in agreement with those obtained by Mayes in his studies. Therefore, if we were to average up the scores we would be able to find the following sequence in order of increasing perseveration.

1. Manic
2. Dementia praecox paranoid
3. Schizophrenic and simple types of dementia praecox
4. Manic-depressive
5. Catatonics

1. Stephenson, "Studies in Experimental Psychology II. Some Comments on Factor Analysis," Journal of Mental Science, 1932, 78, 315.
 2. I. J. Mayes, "Studies in Catatonia: I. Perseverational Tendencies in Catatonia," The Psychiatric Quarterly, 1934, 8, 728.
 3. Stephenson, "Studies in Experimental Psychology II. Some Comments on Factor Analysis," Journal of Mental Science, 1932, 78, 315.

B. Interrcorrelation between the Tests

It is no secret that no battery of perseveration tests is near to being standardized. These tests, which have been previously used by other investigators⁵, do not appear to intercorrelate well for psychotic subjects. Some better results may be obtained with normal subjects whose greater stability and lesser variability may account for it.⁶ The lack of significant correlations between the tests in psychotic subjects does not cast any reflections upon the validity and reliability of the tests. The trouble may be found in the psychotic subject who vacillates and changes from moment to moment though preserving general tendencies like the manic "low" and the depressive "high" perseveration.

The results also make apparent a significant connection between the two sensory tests and a less significant one between the two motor tests and little or practically no connection between the two types of tests (sensory and motor). Perhaps Allport's objection in this connection is pertinent (regarding the sensory tests).⁷ "....they are too trivial to tap the developed volitional functions of personality." Though in view of the definition of perseveration (in the broad sense) it does not necessarily have to meet with Allport's qualifications. Perseveration is not necessarily part of the "volitional functions of personality." The tendency to continue an activity can be well understood without bringing in the possible facilitating or inhibiting effects of volition.

The above discussion of facts does not necessarily destroy the value

⁵ See above; Chapter II.

⁶ B.J. Shevach, "Studies in Perseveration VII. Experimental Results of Tests for Sensory Perseveration," The Journal of Psychology, 1936, 3, 403-427.
⁷ G.W. Allport, Personality, A Psychological Interpretation, 1937.

Interrelationships between the Tests

It is no secret that no battery of personality tests is new to the world. These tests which have been previously used by other investigators, do not appear to intercorrelate well for psychotic subjects. A better results may be obtained with normal subjects whose general ability and hence variability may account for it. The lack of significant correlations between the tests in psychotic subjects does not mean reflections upon the validity and reliability of the tests. The results may be found in the psychotic subject who is unstable and changes a moment to moment though generally general tendencies like the manic and the depressive "high" pervasiveness.

The results also make apparent a significant connection between the sensory tests and a less significant one between the two motor tests and the or practically no connection between the two types of tests (sensory and motor). Perhaps Allport's objection in this connection is pertinent regarding the sensory tests. "...they are too trivial to tap the volitional functions of personality." Though in view of the function of pervasiveness (in the broad sense) it does not necessarily follow that with Allport's qualification, pervasiveness is not necessarily of the "volitional functions of personality." The tendency to continue activity can be well understood without bringing in the possible inhibition or inhibiting effects of volition.

The above discussion of facts does not necessarily destroy the value

¹ See above Chapter II.
² E. S. Shaffer, "Studies in Pervasive Personality," Experimental Results of the Sensory Perversion, The Journal of Psychology, 1936, 3, 403-43.
³ C. E. Allport, Personality: A Psychological Interpretation, 1937.

of the results obtained and individual differences uncovered in the psychotic groups.

The test on "Rhythmic Perseveration" shows that the subjects who did not change their rhythm at all, are also higher perseverators on the other tests. However, on the basis of this dichotomous classification of perseverators and non-perseverators it is difficult to work out a correlational function. The greater tendency of correspondence with this dichotomy is on part of the motor tests while the results of the sensory tests are equivocal. It must be concluded that this measure of perseveration has great promise; it may best tap the true sources of the perseverative tendency after greater refinement is introduced in the recording apparatus.

C. Relation between Age and Perseveration

In our group comparisons we tried to eliminate the age factor since some of the tests correlate positively with age. The Adaptation experiment results and those of electrical sensitivity and the Z-test correlate positively and quite consistently (preliminary and final results) with age. Granted that some of the correlations are not very significant; (low correlation and high probable error) they indicate a definite tendency nevertheless. On the whole, our results are in agreement with those of Cattell⁸ obtained from normal subjects with motor tests only. There is a lower perseveration score in the early twenties than in later years. The low ratios of the differences between the means and the probable errors of those differences (less than 4) may detract from the absolute certainty,

⁸ R.B.Cattell, "On the Measurement of 'Perseveration,'" British Journal of Educational Psychology, 1935, 5, 76-92.

that a true difference exists. With the exception of the stroke test the tendency for the "p" score to increase with age (until 50) is quite marked. However, we were unable to confirm and must disagree with his conclusion that there is "....a further rise in extreme old age."⁹ Of course, it also depends on the definition of "extreme old age." But on the contrary, our results show a slight decrease in the "p" scores when the subjects approach sixty. This objection, however, must be qualified and it must be understood that complications due to psychosis are a factor which cannot be overlooked in the comparison. But, for the sake of truth, it should be stated that the differences between the means in Cattell's work are rather small; also, no measures of significance are given, which robs the work of its conclusiveness.

Another explanatory observation must be made before closing this section. The "lower p" score in motor tests only may be also explained on the basis of a reduction in general speed and fluency with age. Since, to some extent, the "p-score" depends on the initial speed of performance, it is accordingly reduced at the older age.

D. "Improvement" and P-Scores

Stephenson's¹⁰ claims that scores on perseveration tests may serve as a "barometer" of the mental state of the patient, are somewhat exaggerated and unwarranted by the evidence. Our results do show a tendency toward lower "p" scores in the case of improved patients. The degree of improvement of the patients in our results is, however, determined on more objective

⁹ ibid., 90.

¹⁰ Stephenson, op.cit.

53

...a first difference exists. With the exception of the stroke test the tendency for the "p" score to increase with age (until 50) is quite marked. However, we were unable to confirm and not disagree with his conclusion. There is "...a further rise in extreme old age." Of course, it also ends on the definition of "extreme old age." But on the contrary, our data show a slight decrease in the "p" scores when the subject approach 50. This objection, however, must be qualified and it must be understood that complications due to psychosis are a factor which cannot be looked in the comparison. But, for the sake of truth, it should be noted that the differences between the means in Cattell's work are rather small; also, no measures of significance are given, which robs the sort of conclusiveness.

Another explanatory observation must be made before closing this section. The "lower p" found in motor tests only may be also explained on basis of a reduction in general speed and fluency with age. Since, to a certain extent, the "p-score" depends on the initial speed of performance, it is naturally reduced in the older age.

Personality and P-Scores

It is known that a high "p" score on Cattell's tests may serve as a "personality" of the mental state of the patient, and somewhat exaggeratedly interpreted by the evidence. Our results do show a tendency toward a "p" score in the case of improved patients. The degree of improvement in patients in our results is, however, determined on more objective

bases. Also, Stephenson retested patients who improved and based his conclusions on the results obtained from them. This procedure is not quite valid, since the element of training is not taken into consideration.

Darroch¹¹ investigated this point and found that it may be a vitiating factor. Our comparisons are not based on retests. We did not administer the same tests to the same subjects during different stages of recovery. Our results are based on a comparison of scores of groups of subjects tested only once. These groups were differentiated, as stated above, on the basis of the degree of improvement. Hence our results seem to be more reliable.

E. Theoretical Considerations.

On the basis of our results, we cannot agree with most of the investigators who put dementia praecox as highest in regard to perseveration. As we pointed out, there was no sufficient differentiation between the groups within the general group in their investigations.

The results, however, which follow from the work of Stephenson and others, seem to possess theoretical implications which are in keeping with some traditional typological theories. It was mentioned earlier in this paper¹³ that Gross identified the low secondary function with the "broad and shallow" type of consciousness while a high secondary function was characteristic of the "deep and narrow" type of consciousness. Jung's Extroversion-Introversion is parallel to Gross' types. Jung says¹⁴ that

¹¹ J. Darroch, "An Investigation into the Degree of Variation in the Score of a Motor Perseveration Test," British Journal of Psychology, 1938, 31, 248-262.

¹² Ibid., 261.

¹³ Gross, op.cit.

¹⁴ C. C. Jung, Psychological Types, 1926.

... also, depression reported patients who improved and based the
 conclusions on the results obtained from them. This procedure is not valid
 at all, since the element of training is not taken into consideration.
 If I investigated this point and found that it may be a violating
 factor, the comparisons are not based on reliability. We did not administer
 a same test to the same subjects during different stages of recovery.
 The results are based on a comparison of scores of groups of subjects
 at only one stage. These groups were differentiated, as stated above,
 on the basis of the degree of improvement. Hence our results seem to be
 reliable.

Theoretical Considerations

On the basis of our results, we cannot agree with most of the
 investigators who put dementia praecox as highest in regard to perseveration.
 We pointed out, there was no sufficient differentiation between the
 cases within the general group in their investigations.
 The results, however, which follow from the work of Stephenson and
 others, seem to possess theoretical implications which are in keeping with
 a traditional hypothetical theories. It was mentioned earlier in this
 paper that Gross identified the low secondary function with the "gross"
 shallow type of consciousness while a high secondary function was
 characteristic of the "deep and narrow" type of consciousness. Lurie's
 perseveration-introversion is parallel to Gross' types. Lurie says that

... J. B. Watson, "An investigation into the Degree of Variation in the
 of a Motor Perseveration Test," British Journal of Psychology, 1925, 2

12, 1925, 221.
 13 Gross, op. cit.
 14 Lurie, op. cit.

"Introversion is distinguished by....intensive primary function and a correspondingly long secondary function," while "extroversion is characterized by weak primary function, and a correspondingly short secondary function." Since dementia praecox is identified with extreme introversion, and manic-depression with extreme extroversion, the two psychotic groups should give, deducing from Jung, high and low perseveration scores, respectively. Our results, as stated above, do not "support this notion." We are opposed to the broad classification, in regard to perseveration, of manic-depressives and dementia praecox. We find that in both classes there are groups of individuals who have either high or low perseveration. This is not due to any "supervening dementia," but to the peculiar state of the individual personality. Jung admits that "exceptions, however, are frequent even in one and the same individual."¹⁵ An introvert may be extroverted under favorable conditions, while an extrovert may turn inward in a hostile, unfriendly situation.

We cannot, then, always speak of the manic-depressive, extrovert constitution, neither can we speak of the dementia praecox, introvert constitution. It appears that the psychiatric condition itself, ultimately, perhaps, traced to psychological extremes of types, creates the new milieu for the individual and guides and changes the secondary function or perseverative tendency along with it.

Thus, the manic, whose milieu is "favorable", is extremely extroverted, while the depressive of, presumably, the same constitution, gets "introverted" as a result of his own unpleasant world. This can also similarly be traced in the dementia praecox cases, who also become extroverted, while

¹⁵ Ibid., 356.

innovation is distinguished by... intensive primary function and a
 correspondingly long secondary function, while "introversion is character-
 ized by... weak primary function, and a correspondingly short secondary
 function." Since dementia praecox is identified with extreme introversion,
 manic-depression with extreme extroversion, the two psychotic groups
 can give, deducting from 100, high and low perversion scores,
 respectively. Our results, as stated above, do not support this notion.
 are opposed to the broad classification, in regard to perversion, of
 manic-depressives and dementia praecox. We find that in both classes there
 are groups of individuals who have either high or low perversion. This
 can be due to any "supervening dementia," but to the peculiar state of the
 individual personality. There is no doubt that "exceptional, however, are frequent
 in one and the same individual."¹² An introvert may be extroverted
 in favorable conditions, while an extrovert may turn inward in a hostile,
 friendly situation.
 We cannot, then, always speak of the manic-depressive, extrovert
 or the dementia praecox, introvert. Neither can we speak of the dementia praecox, introvert
 or extrovert. It appears that the psychiatric condition itself, ultimately,
 leads, through to psychological extremes of types, creates the new milieu
 for the individual and guides and changes the secondary function of perva-
 sion tendency along with it.
 Thus, the manic, whose milieu is "extroverted," is extremely extroverted,
 and the depressive, whose milieu is "introverted," is extremely intro-
 verted. This can also similarly
 be seen in the dementia praecox cases, who also become extroverted, while

the catatonics of that group are "true to their classification" as extreme introverts and perseverators.

More caution, then, will have to be exercised in our linking extroversion and introversion with high and low perseveration. The particular, though transient, state of the individual under the experimental conditions must be considered. The emotional tone is an important factor. And, perhaps, more conclusive evidence may substantiate Shevach's¹⁶ statement, that "the subjects must first be divided into those in whom perseveration is a functional unit and those in whom it is highly specific."

We must then, from our work with psychotics, conclude that extremes of extroversion and of introversion go with extremes of low and high secondary function. However, the secondary function is also changeable in correspondence with the changeability of the state of introversion and extroversion.

Returning to Shevach's statement quoted above, we may consider it as a reasonable explanation of the low intercorrelation between our perseveration tests. A number of our individuals may be perseverators "true and consistent," while in others, perseveration does not exist as a functional unit, but is "highly specific" in the various and sundry test-situations.

¹⁶ B.J. Shevach, "Studies in Perseveration VII. Experimental Results of Tests for Sensory Perseveration," The Journal of Psychology, 1936, 3, 403-427.

...of that group are "true to their classification" as extreme
verts and perseverators.
...more careful, then, will have to be exercised in our linking extro-
...and introversion with high and low perseveration. The particular
...of the individual under the experimental conditions
...be considered. The emotional tone is an important factor. And,
...more conclusive evidence may substantiate Shewach's statement,
...the subjects must first be divided into those in whom perseveration is
...functional unit and those in whom it is highly specific."
We must then, from our work with psychotics, conclude that extreme
...of introversion go with extremes of low and high
...function. However, the secondary function is also changeable in
...with the changeability of the state of introversion and
...version.
Returning to Shewach's statement quoted above, we may consider it as
...possible explanation of the low introversion between our persev-
...tests. A number of our individuals may be perseverators "true and
...," while in others, perseveration does not exist as a functional
...but is "highly specific" in the various and sundry test-situations.
16. J. Shewach, "Studies in Perseveration VII. Experimental Results
...for Sensory Perseveration," The Journal of Psychology, 1956, 5,
-137.

2. Conclusions

In drawing our conclusions we must be cautious to distinguish between actual factual results obtained in experiments and theoretical super-structures. The mixture of these two elements has been the weakness of the line of argumentation between them, may be a very tempting urge. We shall, however, try to avoid these pitfalls and confine ourselves to the following conclusions warranted by the experimental data:

CHAPTER VII

1. Differences between the two general psychological categories of dementia praecox and schizophrenia, with regard to preservation of memory, are not statistically significant.

CONCLUSIONS

2. Differences between the two general psychological categories of dementia praecox and schizophrenia, with regard to preservation of memory, are not statistically significant.

3. Differences between the two general psychological categories of dementia praecox and schizophrenia, with regard to preservation of memory, are not statistically significant.

4. Differences between the two general psychological categories of dementia praecox and schizophrenia, with regard to preservation of memory, are not statistically significant.

5. The results tend to be lower preservation than are the dementia praecox group.

6. The results tend to be lower preservation than are the dementia praecox group, while the dementia praecox group is the highest in this group.

7. When all groups are compared together, the dementia praecox group is the lowest in preservation, while the dementia praecox group is the highest in preservation.

8. Age differences in dementia praecox have introduced a factor of error in previous investigations. Patients in the third decade of life (20-30 group) have a considerably lower preservation score than the dementia praecox group. There also seems to be a slight tendency for

CHAPTER VII

CONCLUSIONS

A. Conclusions

In drawing our conclusions we must be cautious to distinguish between actual factual results obtained in experimentation and theoretical superstructures. The mixture of these two elements and the eradication of the line of demarcation between them, may be a very tempting urge. We shall, however, try to avoid these pitfalls and confine ourselves to the following conclusions warranted by the experimental data:

1. Differences between the two general psychiatric categories of dementia praecox and manic-depressive psychoses, with regard to perseveration as measured by our tests, exist, but are not sufficiently significant statistically. The tendency is in the direction of higher perseveration in the manic-depressive group and lower in the dementia praecox group.

2. Differences in degree of perseveration also exist between the sub-groups, within each general psychiatric category:

- a. The manics tend to be lower perseverators than are the depressives.

- b. The catatonics are the greatest perseverators in the dementia praecox group, while the paranooids tend to be lowest in this group.

- c. When all groups are compared together, the manics tend to be the lowest perseverators, while the catatonics possess the highest degree of perseveration.

3. Age differences do exist in perseveration and have introduced a factor of error in previous investigations. Patients in the third decade of life (20-29 group) have a considerably lower perseveration score than the individuals of an older age. There also seems to be a slight tendency for

Conclusions

In drawing our conclusions we must be cautious to distinguish between the results obtained in experimental and theoretical studies. The nature of these two elements and the explanation of the difference between them may be a very complex one. We shall, however, try to avoid these pitfalls and confine ourselves to the following conclusions suggested by the experimental data:

1. Differences between the two general psychiatric categories of manic-depressive and manic-depressive psychosis, with regard to perseveration, are suggested by our tests, exist, but are not sufficiently significant statistically. The tendency is in the direction of higher perseveration in the manic-depressive group and lower in the manic-depressive group.

2. Differences in degree of perseveration also exist between the groups, within each general psychiatric category:

- a. The manic tend to be lower perseverators than the depressive.
- b. The manic group has the greatest perseverators in the manic-depressive group, while the depressive tend to be lowest in this group.
- c. When all groups are compared together, the manic tend to be the lowest perseverators, while the depressive possess the highest degree of perseveration.

3. Age differences do exist in perseveration and have introduced a factor of error in previous investigations. Patients in the third decade of life (30-39 group) have a considerably lower perseveration score than the patients of an older age. There also seems to be a slight tendency for

the "p" score to go down after the individual has reached 50 years of age.

4. The tests employed do not seem to give to perseveration the status of functional unity; from our data we are unable to state whether the subjects or the tests were responsible for that.

5. The new test of "rhythmic perseveration" gives some measure of validation of the results of the other tests.

6. The closer proximity of the patient to "normalcy" is also characterized by a lower perseverative tendency.

B. On the Future

We do not claim too much for these results. The contacts of perseveration with psychiatry are numerous and provide a very fertile field for further investigation. Conquest of new frontiers in this respect will be largely conditioned by the perfection of our instrumental and psychometric devices in measuring the perseverative tendency or secondary function. Like many investigators, we attempt to add our bit towards "a complete solution"-yet to be achieved.

1. The facts employed do not seem to give so representative the

of functional unity; from our data we are unable to state whether

subjects of the tests were responsible for what.

2. The test of "rhythmic perseveration" gives some measure of

duration of the results of the other tests.

3. The closer proximity of the patient to "analog" is also

indicated by a lower perseverative tendency.

On the Future

We do not claim too much for these results. The benefits of psycho-

logical with psychology are numerous and growing a very real field for

an investigation. Comparison of new theories in this respect will be

self conditioned by the perfection of our instrumental and psychological

and in measuring the perseverative tendency or secondary function.

many investigators, we attempt to add our bit towards the complete

achievement to be achieved.

BIBLIOGRAPHY

- Allport, G. W., Personality, A Psychological Interpretation. New York: Henry Holt and Company, pp. 196, 244, 416-418, 1937.
- Bernreuter, R. G., Manual for the Personality Inventory. Stanford University Press, 1935.
- Bernstein, G., "Quickness and Intelligence." British Journal of Psychology, Mon. Suppl., 1924, VII.
- Best, C. H. and Taylor, N. B., The Physiological Basis of Medical Practice. Baltimore: William Wood and Company, 1937.
- Bieshevrel, S., "The Measurement of the Threshold for Flicker and its Value as a Perseveration Test." The British Journal of Psychology, 1938,
- Bleuler, E., Lehrbuch der Psychiatrie. Berlin: Verlag von Julius Springer, 43, 56, 62, 110, 61, 1918.
- Britt, S. H., "Theories of Retroactive Inhibition." Psychological Review, 1936, 43, 207.
- Boring, E. G., A History of Experimental Psychology. (3rd. printing) New York: The Century Company, 1933.
- Cameron, D. E., Objective, Experimental Psychiatry. New York: The MacMillan Company, 16, 70-73, 1935.
- _____, "Mensuration in the Psychoses." American Journal of Psychiatry, 1933, XIII, 153-169.
- Cameron, D. E. and Caunt, T. G. B., "Studies in Perseveration." Journal Mental Science, 1933, 79, 735-745.
- Cathcart, E. P. and Dawson, S. I., "Persistence: A Characteristic of Remembering." British Journal of Psychology, 1928, 18, 262-75.
- Cattell, R. B., "On the Measurement of 'Perseveration.'" British Journal of Educational Psychology, 1935, 5, 76-92.
- _____, "Perseveration and Personality---Some Experiments and a Hypothesis." Journal of Mental Science, 1935, 81, 151-167.
- Crutcher, R., "An Experimental Study of Persistence." Journal Applied Psychology, 1934, 18, 409-417.

References

- Adams, G. E., Personality: A Psychological Approach. New York: Henry Holt and Company, 1937, 2nd ed., 1957.
- Adams, G. E., Manual for the Personality Inventory. Evanston: University Press, 1953.
- Adams, G. E., "Consciousness and Intelligence," British Journal of Psychology, 1954, 46, 171.
- Adams, G. E. and Taylor, K. H., The Psychological Basis of Mental Illness. Baltimore: William Wood and Company, 1957.
- Adams, G. E., "The Measurement of the Threshold for Sleep and the Sleep as a Performance Test," The British Journal of Psychology, 1955, 46, 171.
- Adams, G. E., Handbook for Psychological Testing: Volume 1: The Student. New York: Holt, Rinehart and Company, 1957, 2nd ed., 1957.
- Adams, G. E., "Theories of Retrospective Amnesia," Psychological Review, 1957, 64, 207.
- Adams, G. E., A History of Experimental Psychology. (2nd. printing) New York: The Century Company, 1955.
- Adams, G. E., Introductory Psychology. New York: The Century Company, 1955, 10-12, 1955.
- Adams, G. E., "Amnesia in the Laboratory," American Journal of Psychology, 1955, 68, 121-122.
- Adams, G. E. and Gault, E. H., "Amnesia in the Laboratory," Journal of Experimental Psychology, 1955, 50, 755-756.
- Adams, G. E. and Gault, E. H., "Amnesia in the Laboratory," Journal of Experimental Psychology, 1955, 50, 755-756.
- Adams, G. E., "The Measurement of Performance," British Journal of Psychology, 1955, 46, 171.
- Adams, G. E., "The Measurement of Performance," British Journal of Psychology, 1955, 46, 171.
- Adams, G. E., "The Measurement of Performance," British Journal of Psychology, 1955, 46, 171.
- Adams, G. E., "The Measurement of Performance," British Journal of Psychology, 1955, 46, 171.
- Adams, G. E., "The Measurement of Performance," British Journal of Psychology, 1955, 46, 171.

- Cushing, H. M., "A Perseverative Tendency in Pre-School Children." Archives of Psychology, 1929, 106.
- Darroch, J., "An Investigation into the Degree of Variation in the Score of a Motor Perseveration Test." British Journal of Psychology, 1938, 28, 248-262.
- Dorcus, R. M., "Is there a 'Unit-Trait' of Volitional Perseveration?" Journal of General Psychology, 1935, 13, 345-356.
- Dybowski, Mieczysław, "Perseveration as a Measure of Negativism, and its Estimation by Means of Photographs." Kwartalnik Psychologiczny, 1935, VI, 221-236.
- Edwards, K. H. R., "The Perseverative Tendency," The Journal of Abnormal and Social Psychology, 1933, 28, 198-203.
- Eisenson, J., "A Note on the Perseverating Tendency in Stutterers." Journal of Genetic Psychology, 1937, 50, 195-8.
- Ewen, John H., "Perseveration in the Insane Epileptic." Journal of Mental Science, 1930, 76, 537-540.
- Foster, W. S., "On the Perseverative Tendency." American Journal of Psychology, 1914, 25, 393-428.
- Freud, S., Beyond the Pleasure Principle. The International Psychoanalytic Press, London, p. 44, 1922.
- Garrett, H. E., Statistics in Psychology and Education. Longmans, Green and Company: New York, 1937.
- Gross, O., Die Zerebrale Sekundaerfunktion. Leipzig, p. 69, 1902.
- Guilford, J. P., Psychometric Methods. McGraw Hill Book Company: New York, 1936.
- Guirdham, Arthur, "Rorschach Test in Epileptics." Journal of Mental Science, 1938, 84, 887-892.
- Hargreaves, H. L., "The Faculty of Imagination." British Journal of Psychology, monograph Supplement, 1927, 10, 1-74.
- Henderson, D. K. and Gillespie, R. D., A Textbook of Psychiatry, (3rd. ed.) pp. 83, 86, 210, 1933.
- Hecht, S., "The Nature of Foveal Dark Adaptation." Journal of General Physiology, 1919-20, 2, 499.
- Heymans, C. and Brugmans, H., "Intelligenzprüfungen mit Studierenden." Zeitsch. f. Psych., 1913, 7, 317-331.

- Hunt, J. McV. and Guilford, J. P., "Fluctuation of an Ambiguous Figure in Dementia Praecox and Manic Depressive Patients." The Journal of Abnormal and Social Psychology, 1932-33, 27, 443-452.
- Jasper, H. H., "Is Perseveration a Functional Unit Participating in all Behavior Processes." Journal of Social Psychology, 1931, 2, 28-51.
- Jones, L. Wynn, "An Investigation into the Significance of Perseveration." Journal of Mental Science, 1926, 74, 653-659.
- Jung, C. G., Collected Papers on Analytical Psychology, Bailliere, Tindal and Company: London, pp. 106, 111, 1922.
- _____, Psychological Types, Harcourt, Brace and Company, pp. 336, 357, 1926.
- Kendig, I. and Shevach, B. J., "Studies in Perseveration I. A Survey of Researches in Perseveration." The Journal of Psychology, 1936, 3, 223-230
- Lankes, W., "Perseveration." British Journal of Psychology, 1915, 7, 367-419.
- Mays, L. L., "Studies in Catatonia V. Perseverational Tendencies in Catatonic Patients." The Psychiatric Quarterly, 1934, 6, 726-730.
- McGeoch, John A., "Re: Retroaction and Perseveration." American Journal of Psychology, 1933, 45, 537.
- Muller, G. E. and Pilzecker, H. J., "Studien der Gedächtniss." Zsch. f. Psychol., 1900, Erg. 1.
- Neisser, C., "65 Sitzung Des Vereins Ostdeutscher Irren Aerzte." Allg. Zsch. f. Psychiatrie, 1894, 51, 1016.
- Pavlov, I. P., "An Attempt at a Physiological Interpretation of Obsessional Neurosis and Paranoia." Journal of Mental Science, 1934, 80, 187-197.
- Pinard, J. W., "Tests of Perseveration." British Journal of Psychology, 1932, 23, 5-19.
- _____, "The Relation to Certain Psychopathic Conditions and to Introversion." British Journal of Psychology, 1932, 23, 114-127.
- Remington, G. A., Gleanings of Low Voltage Techniques. 1930.
- Rogers, K. H., "Perseveration." Journal of Mental Science, 1935, 81, 138-44.
- _____, "Perseveration and Personality." Journal of Mental Science, 1935, 81, 145-150.
- _____, "A Relationship Between Perseveration and School Achievement." Psychological Bulletin, 1930, 30, 663-669.

V. W. and G. W. (1937). "The Psychology of an Antisocial Type in
 Domestic Violence and Mental Disorders." The Journal of
 Abnormal and Social Psychology, 1937, 32, 421-432.

H. H. (1937). "The Psychology of a Psychotic Type of
 Behavior." Journal of Social Psychology, 1937, 32, 78-81.

E. W. (1937). "An Investigation into the Psychology of 'Parasitism'." Journal of Social Psychology, 1937, 32, 82-85.

C. C. (1937). Collected Papers on Psychological Problems. London: Tavistock
 and Company, Ltd., 1937.

Psychological Types. Harcourt, Brace and Company, 1937, 1937.

J. and S. (1937). "Studies in the Psychology of a Survey of
 Research in Psychology." The Journal of Psychology, 1937, 32,
 225-230.

W. (1937). "Parasitism." British Journal of Psychology, 1937, 32, 237-
 242.

E. L. (1937). "Studies in the Psychology of Parasitism in
 Domestic Violence." The Journal of Psychology, 1937, 32, 238-250.

J. (1937). "The Psychology of Parasitism." American Journal
 of Psychology, 1937, 32, 251.

E. L. and E. W. (1937). "Studies in the Psychology of Parasitism." Journal
 of Psychology, 1937, 32, 252.

C. C. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 253-254.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 255-256.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 257-258.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 259-260.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 261-262.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 263-264.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 265-266.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 267-268.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 269-270.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 271-272.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 273-274.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 275-276.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 277-278.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 279-280.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 281-282.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 283-284.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 285-286.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 287-288.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 289-290.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 291-292.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 293-294.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 295-296.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 297-298.

J. (1937). "The Psychology of Parasitism." Journal of Psychology, 1937, 32,
 299-300.

Shevach, B. J., A Study of Sensory Perseveration, submitted to the Department of Philosophy and Psychology of Harvard University in partial fulfillment of the requirements for the degree of Doctor of Philosophy, April, 1935.

_____, "Studies in Perseveration VI. Methods for the Study of Sensory Perseveration." The Journal of Psychology, 1936, 3, 381-402.

_____, "Studies in Perseveration VII. Experimental Results of Tests for Sensory Perseveration." The Journal of Psychology, 1936, 3, 403-427.

Shipley, Walter C., "Studies of Catatonia VI. Further Investigation of the Perseverative Tendency." Psychiatric Quarterly, 1934, 8, 736-744.

Slocombe, C. S. and Brakeman, E. E., "Psychological Tests and Accident Proneness." British Journal of Psychology, 1930, 21, 29-38.

Smith, Carl E., "The Effect of Changes in the Temperature and Humidity of the Air on the Apparent Skin Resistance." The Journal of Psychology, 1937, 3, 325-331.

Spearman, C., The Abilities of Man. New York: The MacMillan Company, pp. 42, 51, 291-307, 356, 391, 404, 412, 1927.

_____, "The Tenth Maudsley Lecture." Journal of Mental Science, 1929, 75, 357-370.

Stephenson, W., "Studies in Experimental Psychiatry I. A Case in General Inertia." Journal of Mental Science, 1931, 77, 723-741.

Stephenson, W., "Studies in Experimental Psychiatry II. Some Contacts of p-factor with Psychiatry." Journal of Mental Science, 1932, 78, 315-330.

_____, "Studies in Experimental Psychiatry: III. p-score and Inhibition for High p-Praecox Cases." Journal of Mental Science, 1932, 78, 908-920.

_____, "Perseveration and Character." Character and Personality, 1935, 4, 44-52.

_____, "Studies in Perseveration." Nature, 1934, 134, 860.

_____, "An Introduction to So-Called Motor Perseveration Tests." British Journal of Educational Psychology, 1934, IV, 186-207.

Travis, L. E. and Knott, R. J., "Brain Potential Studies of Perseveration I. Perseveration Time to Light." Journal of Psychology, 1936, 3, 97-100.

Wash, B. J., A Study of Human Personality, submitted to the
Department of Psychology and Psychiatry at Harvard University as
partial fulfillment of the requirements for the degree of Doctor
of Philosophy, April 1935.

_____, "Studies in Personality VI. Studies for the Study of
Personality I. Personality," The Journal of Psychology, 1935, 3, 451-463.

_____, "Studies in Personality VII. Experimental Studies of
Personality I. Personality," The Journal of Psychology, 1935,
3, 463-477.

_____, "Studies in Personality VIII. Further Investigation of the
Personality I. Personality," The Journal of Psychology, 1935, 3, 477-489.

_____, "Studies in Personality IX. The Psychological Tests and Methods
I. Personality," The Journal of Psychology, 1935, 3, 489-501.

_____, "Studies in Personality X. The Nature and Meaning of
the Personality I. Personality," The Journal of Psychology,
1935, 3, 501-511.

_____, "Studies in Personality XI. The Personality I. Personality,"
The Journal of Psychology, 1935, 3, 511-521.

_____, "Studies in Personality XII. The Personality I. Personality,"
The Journal of Psychology, 1935, 3, 521-531.

_____, "Studies in Personality XIII. The Personality I. Personality,"
The Journal of Psychology, 1935, 3, 531-541.

_____, "Studies in Personality XIV. The Personality I. Personality,"
The Journal of Psychology, 1935, 3, 541-551.

_____, "Studies in Personality XV. The Personality I. Personality,"
The Journal of Psychology, 1935, 3, 551-561.

_____, "Studies in Personality XVI. The Personality I. Personality,"
The Journal of Psychology, 1935, 3, 561-571.

_____, "Studies in Personality XVII. The Personality I. Personality,"
The Journal of Psychology, 1935, 3, 571-581.

_____, "Studies in Personality XVIII. The Personality I. Personality,"
The Journal of Psychology, 1935, 3, 581-591.

_____, "Studies in Personality XIX. The Personality I. Personality,"
The Journal of Psychology, 1935, 3, 591-601.

Travis, L. E. and Knott, R. J., "Brain Potential Studies of Perseveration
11. Perseveration Time to Visually Presented Words." Journal of
Experimental Psychology, 1937, 21, 353.

Troland, L. T., The Fundamentals of Human Motivation. New York: D. Van
Nostrand Company, p. 395, 1928.

Viteles, Morris S., Industrial Psychology. New York: W. W. Norton and
Company, 1932.

White, W. A., Outlines of Psychiatry. Washington: Nervous Mental Diseases
Publishing Company, 13th ed., pp. 103, 252, 1932.

Wiersma, E., "Die Sekuendaerfuenktion bei Psychosen." Journal f. Psychol.
u. Neurologie, 1906, VIII, 1-24.

_____, "Psychology of Dementia." Journal of Mental Science, 1930, 76.

Yule, E. P., "The Resemblance of Twins with Regard to Perseveration."
Journal of Mental Science, 1935, 81, 489-501.

10. L. E. and Knott, H. J., "Visual-Motor Studies of Retention
 11. Retention Time for Verbal Material." Journal of
Experimental Psychology, 1937, 21, 553.

12. L. E., The Fundamentals of Human Psychology, New York: N. Van
 Nostrand Company, 1937, 1937.

13. Knott, H. J., Industrial Psychology, New York: N. Van
 Nostrand Company, 1937.

14. W. A., Studies of Individual Differences, New York: World
 Publishing Company, 1937, 1937, 1937.

15. E., "The Correlation of Retention and Intelligence," Journal of
Experimental Psychology, 1937, 21, 1-21.

16. "Psychology of Learning," Journal of Mental Science, 1930, 75.

17. E. P., "The Relationship of Verbal and Visual Retention,"
Journal of Mental Science, 1937, 82, 415-422.



The candidate was born in the town of Merkini, Lithuania, on June 20, 1912. He is the oldest son of David J. Rabin and Sara Simon Rabin. After graduation from the local grammar school, he entered the Kaunas "Gimnasium" from which he graduated in 1929. The regular curriculum included a great deal of mathematics, physics and humanitarian studies. Hebrew, Lithuanian and German were thoroughly studied, while Russian and Polish were less important in the candidate's education.

In the year of 1930, upon his arrival to the United States, the candidate entered Northeastern University as a candidate for the Bachelor of Science degree in the department of electrical engineering. In 1934 he transferred to the Boston University School of Education, from which he graduated in 1935. The following year (1936) he received the Master of Arts degree in the department of psychology of the Boston University Graduate School. The following three years were spent at the Boston University and Harvard Graduate Schools.

The candidate was appointed fellow in the department of psychology at Boston University for the year 1935-36. He spent the following two years as graduate assistant in the same department. On various occasions

The candidate was born in the town of Meriden, Lithuania, on June 1918. He is the oldest son of David J. Rebin and Sara Simon Rebin. After graduation from the local grammar school, he entered the Lithuanian "Gymnasium" from which he graduated in 1933. The regular curriculum included a great deal of mathematics, physics and naturalistic studies. French, Lithuanian and German were thoroughly studied, while Russian and Polish were less important in the candidate's education.

In the year of 1930, upon his arrival to the United States, the candidate entered Northwestern University as a candidate for the Bachelor of Science degree in the department of electrical engineering. In 1934 transferred to the Boston University School of Education, from which he graduated in 1936. The following year (1936) he received the Master of Science degree in the department of psychology of the Boston University Graduate School. The following three years were spent at the Boston University and Harvard Graduate Schools.

The candidate was associated fellow in the department of psychology Boston University for the years 1935-36. He spent the following two years as graduate assistant in the same department. On various occasions

he lectured to classes at the University and substituted as an instructor in psychology at Emerson College in Boston. For the past two summers he acted as camp director at a private boys' camp. He has been engaged in research at the Boston State Hospital since last fall.

lectured to classes at the University and subordinated as an instructor
psychology at Harvard College in Boston. For the past two summers
acted as camp director at a private boys' camp. He has been engaged in
research at the Boston State Hospital since last fall.

PERSEVERATION IN DEMENTIA PRAECOX AND IN THE MANIC DEPRESSIVE PSYCHOSES.
BOSTON UNIVERSITY

The phenomena of perseveration GRADUATE SCHOOL

be traced back to ancient philosophical literature. It is noticeable in every day life, in the uncontrollable persistence of ideas in our minds and in the after effects of previous activities and patterns of behavior upon our present actions. After sensations, after images and the like have also been included under the term perseveration.

Abstract of the Dissertation

It is not an isolated phenomenon since its existence can be explained

PERSEVERATION IN DEMENTIA PRAECOX AND MANIC DEPRESSIVE PSYCHOSES

spontaneous discharges and "after-discharges" of nerve cells occupy the position of established facts. This automaticity and after-discharge in

by

Israel Rabin

the nervous system, after the original stimuli or stimulating situations

(B.S.in Ed., Boston University, 1935; A.M., Boston University, 1936)

mental inertia or perseveration.

The word "perseveration" as such was used, by the German psychiatrist, Heimerl, for the first time. Since the end of the last century, when the term was introduced, it was employed descriptively in psychiatry. Repetitive movements or words on the part of a mental patient, despite his desire

to change his behavior, submitted in partial fulfilment of the requirements for the degree of

Doctor of Philosophy

stimulus, but the response is a stereotyped activity, which was for a while considered synonymous with perseveration, in an oft-repeated activity or postural attitude which cannot be traced back to any stimulus.

1939

Phenomena of a similar nature were observed by Jung in the association experiments. Certain words were heard repeated several times in response to different stimulus words.

BOSTON UNIVERSITY

GRADUATE SCHOOL

Abstract of the Dissertation

PERSEVERATION IN DEMENTIA PARVOX AND MANIC DEPRESSIVE PSYCHOSES

by

Israel Rubin

S.C. in Ed., Boston University, 1933; A.M., Boston University, 1935

submitted in partial fulfillment of the

requirements for the degree of

Doctor of Philosophy

1933

PERSEVERATION IN DEMENTIA PRAECOX AND IN THE MANIC DEPRESSIVE PSYCHOSES.

The phenomenon of perseveration has been recognized long ago and may be traced back to ancient philosophical literature. It is noticeable in every day life, in the uncontrollable persistence of ideas in our minds and in the after effects of previous activities and patterns of behavior upon our present actions. After sensations, after images and the like have also been included under the term perseveration.

It is not an isolated phenomenon since its existence can be explained on the basis of the latest developments in neurophysiology. Automatic and spontaneous discharges and "after-discharges" of nerve cells occupy the position of established facts. This automaticity and after -discharge in the nervous system, after the original stimuli or stimulating situations are removed, is in accordance with the psychologically noted "after-effect;" mental inertia or perseveration.

The word "perseveration" as such was used, by the German psychiatrist, Neisser, for the first time. Since the end of the last century, when the term was introduced, it was employed descriptively in psychiatry. Repetitive movements or words on the part of a mental patient, despite his desire to change his behavioral pattern, were considered to be demonstrations of the perseverative tendency. This repetition, usually, has an original stimulus, but the response could not be checked, while stereotypy, which was for a while considered synonymous with perseveration, is an oft-repeated activity or postural attitude which cannot be traced back to any stimulus. Phenomena of a similar nature were observed by Jung in his association experiments. Certain words have been repeated several times as responses to different stimulus words.

The phenomenon of perseveration has been recognized long ago and may be traced back to ancient philosophical literature. It is noticeable in everyday life, in the uncontrollable persistence of ideas in our minds and the after effects of previous activities and patterns of behavior upon present actions. After sensations, after images and the like have also been included under the term perseveration.

It is not an isolated phenomenon since its existence can be explained on the basis of the latest developments in neurophysiology. Automatic and spontaneous discharges and "after-discharges" of nerve cells occupy the field of established facts. This automaticity and after-discharge in nervous system, after the original stimuli or stimulating situations removed, is in accordance with the psychologically noted "after-effect"; cal inertia or perseveration.

The word "perseveration" as such was used, by the German psychiatrist, Asper, for the first time. Since the end of the last century, when the word was introduced, it was employed descriptively in psychiatry. Repetitive movements or words on the part of a mental patient, despite his desire to change his behavioral pattern, were considered to be demonstrations of perseverative tendency. This repetition, usually, has an original stimulus, but the response could not be checked, while atelecty, which was a while considered synonymous with perseveration, is an off-repeated activity or postural attitude which cannot be traced back to any stimulus.

Moments of a similar nature were observed by Jung in his association experiments. Certain words have been repeated several times as responses to different stimulus words.

On the basis of accumulated clinical observations, Gross, in his theoretical treatise, posited two main principles of all mental life---the primary and secondary functions. The primary function was direct perception and thought in any form, while the secondary function was the "after-function" or the process set up after the removal of the original stimulus. According to Gross, individuals who have a longer secondary function tend to follow a narrow, but deep path in their mental life; once a process is set up, it continues for a long time. On the other hand, the shallow and broad type of person has a shorter secondary function; new stimuli take more rapidly the place of previous mental processes, since there seems to be less secondary function or perseveration. Jung's typology of introversion and extroversion follows the same general outline. He linked his types with perseveration in a similar manner.

The Dutch and English Schools set out, in the beginning of the present century, to investigate experimentally, this secondary-function (synonymous with perseveration). They had to devise tests as well as to correlate existing psychological and psychiatric categories with classes discoverable by means of those tests.

The testing methods followed the following general lines:

- a. Sensory tests were devised; these tests were to measure the duration of the after-effects of different sense stimulations; i.e., the phenomenon of continuance.
- b. Ideational tests which set up a definite response pattern in the subjects; the persistence of this habituated pattern was then measured by the number of responses and by the amount of interference it would show in the following of new instructions; i.e., fluency and interference.

On the basis of accumulated clinical observations, Gross, in his
practical treatise, posited two main principles of all mental life--the
primary and secondary functions. The primary function was direct perception
thought in any form, while the secondary function was the "after-function"
the process set up after the removal of the original stimulus. According
to Gross, individuals who have a longer secondary function tend to follow a
slow, but deep path in their mental life; once a process is set up, it
lingers for a long time. On the other hand, the shallow and broad type of
person has a shorter secondary function; new stimuli take more rapidly the
place of previous mental processes, since there seems to be less secondary
retention or perseveration. Jung's typology of introversion and extroversion
follows the same general outline. He linked his types with perseveration in
a similar manner.

The Dutch and English Schools set out, in the beginning of the
present century, to investigate experimentally, this secondary-function
phenomenon with perseveration). They had to devise tests as well as to
relate existing psychological and psychiatric categories with classes
coverable by means of those tests.

The testing methods followed the following general lines:

- a. Conatory tests were devised; these tests were to measure the
action of the after-effects of different sensory stimulations; i.e., the
persistence of continuance.
- b. Intentional tests, which set up a definite response pattern in the
test; the persistence of this habituated pattern was then measured by
number of responses and by the amount of interference it would show in
following of new instructions; i.e., fluency and interference.

c. Motor tests in which a certain motor activity was initiated and then changed to a similar one, but not the same; the amount of perseveration being the reduction of the output in the second activity due to the after-effects and the interference of the previous one, i.e., interference.

These tests measured perseveration and, at the same time, operationally defined it; however, the functional unity of the entire group was never definitely established.

Experimental work with mental patients did not yield consistent results. In one case melancholics were considered to be higher perseverators than manics, while in another investigation the results were diametrically opposite. Other investigators have "established" the fact that manics are the lowest perseverators in the psychotic group, while manic-depressives are higher and the dementia praecox cases may be separated into two different groups of high and low perseverators. The factor of age was not investigated in connection with this work. Improvement of Dementia Praecox patients was also shown to affect the perseveration scores in the retests. The factor of training that might render the results erroneous was not taken into consideration.

The undetermined state of affairs in the relationship between psychiatric categories, age and improvement and the degree of perseveration, prompted the present investigation. It was felt that age may play a considerable role in determining the "p" score; it was also thought that high and low "p" in dementia praecox must be explained; the necessity to investigate the relationship between improved patients' perseveration-scores and those of the definitely psychotic ones, revealed itself.

Five tests of perseveration were employed in the present investigation.

c. Motor tests in which a certain motor activity was initiated and changed to a similar one, but not the same; the amount of perseveration and the reduction of the output in the second activity due to the after-effects and the interference of the previous one, i.e., interference. These tests measured perseveration and, at the same time, operation-defined it; however, the functional unity of the entire group was never definitely established.

Experimental work with mental patients did not yield consistent results. In one case melancholics were considered to be higher perseverators than manic, while in another investigation the results were statistically opposite. Other investigators have "established" the fact that manic are the lowest perseverators in the psychotic group, while depressives are higher and the dementia praecox cases may be separated into two different groups of high and low perseverators. The factor of age was not investigated in connection with this work. Improvement of dementia praecox patients was also shown to affect the perseveration scores in the tests. The factor of training that might render the results erroneous was taken into consideration.

The undetermined state of affairs in the relationship between psychiatric categories, age and improvement and the degree of perseveration, prompted the present investigation. It was felt that one may give a considerable role in determining the "p" score; it was also thought that high low "p" in dementia praecox must be explained; the necessity to investigate the relationship between improved patients' perseveration-scores and age of the definitely psychotic ones, revealed itself.

Five tests of perseveration were employed in the present investigation.

Two of the tests were sensory---Adaptation to light and Electrical Threshold Sensitivity; two of the other tests were motor---the Z-tests and the Grouped Strokes tests. The fifth test was devised by the present investigator and was called "A Test of Rhythmic Perseveration." The first four tests were used in various investigations by British and American workers. In the last test the subject performs an activity keeping in time with a slow metronome, the speed of which is suddenly changed, and accordingly, the subject's adjustment time to the new speed is considered to be a measure of perseveration. Some of the subjects demonstrated such a degree of perseveration that their speed remained constant throughout the experiment.

Two groups of patients were used. The preliminary group (A) consisted of thirty-two patients, the second experimental group (B) included eighty patients. All of these subjects belonged to two psychiatric categories---dementia praecox and manic-depressive psychoses. They were all males between the ages of nineteen and sixty.

Part of the experimental procedure was the establishment of a rapport between tester and testee. Modes of response had to be changed for some very depressed patients, while in some cases the results were rendered valueless by lack of full comprehension of the instructions and negativism on the part of certain subjects. Elimination of these results did not introduce unscientific sampling since there was no predominance of a particular diagnosis among the negativistic and unreliable subjects.

The preliminary results showed that the depressive group, which was the younger one, had the higher mean perseveration scores than did the praecox patients. There was also some positive correlation between the

of the tests were sensory--adaptation to light and electrical
threshold sensitivity; two of the other tests were motor--the E-tests and
Grouped Stroop tests. The fifth test was devised by the present
investigator and was called "A Test of Hysterical Perseveration." The first
of tests were used in various investigations by British and American
workers. In the last test the subject performs an activity keeping in time
the slow metronome, the speed of which is suddenly changed, and
consequently, the subject's adjustment time to the new speed is considered
as a measure of perseveration. Some of the subjects demonstrated such
degrees of perseveration that their speed remained constant throughout
the experiment.

Two groups of patients were used. The preliminary group (A)
consisted of thirty-two patients, the second experimental group (B) included
fifty patients. All of these subjects belonged to two psychiatric
categories--dementia praecox and manic-depressive psychoses. They were all
between the ages of nineteen and sixty.
Part of the experimental procedure was the establishment of a report
between tester and testee. Modes of response had to be changed for some
depressed patients, while in some cases the results were rendered
useless by lack of full comprehension of the instructions and negativism
the part of certain subjects. Elimination of these results did not
produce unsatisfactory sampling since there was no predominance of a par-
ticular diagnosis among the negativistic and unreliable subjects.
The preliminary results showed that the depressive group, which was
younger one, had the higher mean perseveration scores than did the
schizophrenic patients. There was also some positive correlation between the

tests and age. The difference, however, was not very significant statistically.

After the final results were computed, the following conclusions were drawn:

1. The depressive group tends to consist of high perseverators than does the group of dementia-*praecox* patients.
2. Within the dementia *praecox* group, catatonics are the highest perseverators while the paranoiacs tend to be lowest.
3. With the exception of one test, the general tendency is to establish the following sequence in order of increasing perseveration: mania, dementia *praecox* (paranoia, hebephrenia and simple), depression and catatonia.
4. The lowest "p" scores are obtained by the youngest age group (twenty to twenty-nine) while the others (thirty to fifty-nine) are higher perseverators.
5. The sensory tests correlate satisfactorily with each other; a similar relation exists between the motor tests; from the data obtained, however, there is little evidence of a correlation between the sensory and the motor tests.
6. Patients of improved psychiatric condition tend to be lower perseverators.

Further work with the second group of patients (B) seemed to corroborate the preliminary results. The dementia *praecox* sub-group investigated more thoroughly and matched so as to eliminate th

tests and age. The difference, however, was not very significant.

statistically.

After the final results were computed, the following conclusions were

drawn:

1. The depressive group tends to consist of high perseverators

than does the group of dementia-praecox patients.

2. Within the dementia-praecox group, catatonics are the highest

perseverators while the paranoics tend to be lowest.

3. With the exception of one test, the general tendency is to

establish the following sequence in order of increasing perseveration:

mania, dementia-praecox (paranoic, hebephrenic and simple), depression and

catatonics.

4. The lowest "P" scores were obtained by the youngest age group

(twenty to twenty-nine) while the others (thirty to fifty-nine) are higher

perseverators.

5. The sensory tests correlate satisfactorily with each other;

a similar relation exists between the motor tests; from this data obtained,

however, there is little evidence of a correlation between the sensory and

the motor tests.

6. Patients of improved psychiatric condition tend to be lower

perseverators.

Further work with the second group of patients (B) seemed to per-


corporate the preliminary results. The dementia-praecox sub-

investigated more thoroughly and matched so as to eliminate the

and did the

ion between the

BOSTON UNIVERSITY



1 1719 02556 8645

